arXiv:1210.4667v1 [math.GT] 17 Oct 2012

LOCAL MOVES ON KNOTS AND PRODUCTS OF KNOTS

LOUIS H. KAUFFMAN AND EIJI OGASA

Abstract. We show several relations between local moves on 1-dimensional knots and those on high dimensional knots related by products of knots.

1. Introduction

We show several relations between local moves on 1-knots and those on high dimensional knots (defined in [19, 21, 23]) related by knot products (defined in [8, 9]).

The local moves on 1-dimensional knots (or links) that we discuss are *crossing-changes* and *pass-moves*, which are introduced in [10] (see §3 of this paper): In the pass-move, two 1-dimensional knots (or links) K_+ and K_- differ only in a 3-ball as shown below.


K_+


K_-


K_+


K_-

E-mail: kauffman@uic.edu ogasa@mail1.meijigakuin.ac.jp

Keywords: local moves on 1-knots, local moves on high dimensional knots, crossing-changes on 1-links, pass-moves on 1-links, products of knots, pass-moves on high dimensional links, twist-moves on high dimensional links, branched cyclic covering spaces.

Local moves on high dimensional knots (or codimension two submanifolds in a high dimensional standard sphere) that we discuss are twist-moves, (p, q)-pass-moves, and ribbon-moves. These are defined and are discussed in [19, 20, 21, 22, 23, 24] (see §4). The (p, q)-pass-moves on n-dimensional knots (or n-dimensional submanifolds in S^{n+2}), where $p+q = n+1$ and $p, q, n \in \mathbb{N}$, are as follows: See §4 for detail. Two n-dimensional knots (or n-dimensional submanifolds in S^{n+2}) K_+ and K_- differ only in the $(n+2)$-ball B^{n+2} as shown in the following figures. (B sometimes denotes B^{n+2}.) $B \cap K_+$ (resp. $B \cap K_-$) is diffeomorphic to the disjoint union of $S^{n-p} \times D^p$ and $S^{p-1} \times D^{n+1-p}$. Take an $(n+1)$-dimensional p-handle (resp. an $(n+1)$-dimensional $(n+1-p)$-handle) which is embedded trivially in B^{n+2} and attached to ∂B^{n+2} trivially. Suppose that $S^{n-p} \times D^p$ (resp. $S^{p-1} \times D^{n+1-p}$) is $\overline{\partial h^p - \partial B^{n+2}}$ (resp. $\overline{\partial h^{n+1-p} - \partial B^{n+2}}$). We pass the handle h^p across h^{n+1-p}as in the following figures, move $S^{n-p} \times D^p$ and $S^{p-1} \times D^{n+1-p}$ together and change K_+ into K_-.

Figure 1.1 is a conceptual figure of a $(p, n+1-p)$-pass-move. Here, h^p_+ denotes h^p in $B^{n+2} \cap K_+$. Figure 1.2, which consists of the two figures (1) (2), is another conceptual figure of a $(p, n+1-p)$-pass-move. Note that $(1, 1)$-pass-moves are same as pass-moves on 1-knots (1-links).


$B \cap K_+$
$B \cap K_-$
$S^{n-p} \times D^p$
$= \overline{\partial h^p_+ - \partial B}$
$S^{p-1} \times D^{n+1-p}$
$= \overline{\partial h^{n+1-p} - \partial B}$

Figure 1.1

A $(p, n+1-p)$-pass-move on an n-dimensional submanifold $\subset S^{n+2}$. Note $B = B^{n+2} = D^{n+2} \subset S^{n+2}$. This figure includes h^p_+ and h^{n+1-p}.


$D^1 = [0,1]$
$S^{n-p} \times D^p$
$S^{p-1} \times D^{n+1-p}$
D^{n+1-p}
D^p
This cube is $B = D^{n+2} = D^1 \times D^p \times D^{n+1-p}$
$B \cap K_+$

Figure 1.2.(1)

A $(p, n+1-p)$**-pass-move on an** n**-dimensional submanifold** $\subset S^{n+2}$**.**
Note $B = B^{n+2} = D^{n+2} \subset S^{n+2}$**.**



$B \cap K_-$

Figure 1.2.(2)

A $(p, n+1-p)$**-pass-move on an** n**-dimensional submanifold** $\subset S^{n+2}$**.**
Note $B = B^{n+2} = D^{n+2} \subset S^{n+2}$**.**

The twist-moves on $(2p+1)$-dimensional knots (or codimension two submanifolds in S^{2p+3}) are as follows, where $p \in \mathbb{N} \cup \{0\}$. See §4 for detail. Two $(2p+1)$-dimensional knots (or codimension two submanifolds in S^{2p+3}) K_+ and K_- differ only in the $(2p+3)$-ball B^{2p+3} as shown in the following figures. (B sometimes denotes B^{n+2}.) $B \cap K_+$ (resp. $B \cap K_-$) is diffeomorphic to $S^p \times D^{p+1}$. Take a $(2p+2)$-dimensional $(p+1)$-handle which is embedded trivially in B^{2p+3} and attached to ∂B^{2p+3} trivially. Suppose that $S^p \times D^{p+1}$ is $\overline{\partial h^{p+1} - \partial B^{2p+3}}$. Note that we can twist the handle h^{p+1} in B once (see §4 for detail), move $S^p \times D^{p+1}$ together and change K_+ into K_-.

Figure 1.3, which consists of the two figures (1) (2), is a conceptual figure of a twist-move. The upper half of Figure 1.4 is another conceptual figure of a twist-move. Compare the upper half of of Figure 1.4 and the lower half. If $p = 0$ (i.e., $n = 2p + 1 = 1$), the left figure in the upper half and that in the lower half are same. That is, if $p = 0$, twist-moves are crossing-changes on 1-links. We review another local-move on high dimensional knots (resp. submanifolds), ribbon-moves in §4.



Figure 1.3.(1)

A twist-move on $(2p+1)$**-dimensional submanifold** $\subset S^{2p+3}$**.**
Note $B = D^{2p+3} \subset S^{2p+3}$**.**



Figure 1.3.(2)

A twist-move on $(2p+1)$-dimensional submanifold $\subset S^{2p+3}$.
Note $B = D^{2p+3} \subset S^{2p+3}$.



If $p = 0$



The pair of two ◌ makes a crossing-change on a 1-dimensional link.

Figure 1.4
A twist-move on a 1-link is a crossing-change on a 1-link.

The knot products (or products of knots) are as follows (see [8, 9]). See §5 for detail.

Let J (resp. K) be an n-(resp. m-)dimensional knots (or codimension two high dimensional submanifolds in S^{n+2} (resp. S^{m+2})) where $n, m \in \mathbb{N} \cup \{0\}$. We define a products of knots (or knot product) of the submanifolds J and K if J or K is a fibred submanifold. See fibered submanifolds for §5. Let $J \otimes K$ denote the knot product of J and K. $J \otimes K$ is defined to be an $(n+m+3)$-submanifold $\subset S^{n+m+5}$.

Suppose that $n = 2n'+1$ and $m = 2m'+1$ $(n', m' \in \mathbb{N})$ and that J (resp. K) is defined by using a polynomial $f : \mathbb{C}^{n'+2} \to \mathbb{C}$ (resp. $f : \mathbb{C}^{m'+2} \to \mathbb{C}$) and the intersection of the unit sphere of $\mathbb{C}^{n'+2}$ (resp. $\mathbb{C}^{m'+2}$) and $f^{-1}(0)$ (resp. $g^{-1}(0)$). See [8, 9, 17]. Then $J \otimes K$ is defined by using $f - g$. Of course, in §5, we define $J \otimes K$ if we cannot define J or K by using such a polynomial and if one or two of n and m are even numbers.

One of our main theorems is the following. If a 1-link K is obtained from a 1-link K' by a crossing-change, then $K\otimes$(the Hopf link) is obtained from $K'\otimes$(the Hopf link) by a twist-move (see Theorem 2.1, 2.2, 2.3).

Other results in this paper are as follows: If a 1-knot K is obtained from a 1-knot K' by a pass-move, then $K\otimes$(the Hopf link) is obtained from $K'\otimes$(the Hopf link) by a $(3,3)$-pass-move (see Theorem 2.5).

Let K and K' be 1-knots. The 1-knot K is pass-move equivalent to the 1-knot K' if and only if $K\otimes$(the Hopf link) is $(3,3)$-pass-move equivalent to $K'\otimes$(the Hopf link) (see Theorem 2.5 and 2.7).

Of course we show more results in other high dimensional cases.

2. MAIN RESULTS

We work in the smooth category. Let $L = (K_1, ..., K_m)$ be an m-component n-(dimensional) submanifold $\subset S^{n+2}$. If $m = 1$ and if L is PL homeomorphic to the standard sphere, then L is called an n-dimensional spherical knot. If each K_i is a spherical knot, then L is called an n-dimensional spherical link. We say that n-submanifolds L and L' are *identical* if there exists an orientation preserving identity map $id : S^{n+2} \to S^{n+2}$ such that $id(L)$=L' and $id|_L : L \to L'$ is an orientation and order preserving identity map. We say that n-submanifolds L and L' are *equivalent* if there exists an orientation preserving diffeomorphism $f : S^{n+2} \to S^{n+2}$ such that $f(L)$=L' and $f|_L : L \to L'$ is an orientation and order preserving diffeomorphism. An m-component n-submanifold $L = \{L_1, ..., L_m\}$ is called a *trivial* $(n$-$)$*link* if each L_i bounds an $(n+1)$-ball B_i trivially embedded in S^{n+2} and if $B_i \cap B_j = \phi (i \neq j)$. If $m = 1$, then L is called a *trivial* $(n$-$)$*knot*.

The following theorems are special cases of our results.

Theorem 2.1. (This follows from Theorem 8.2.) *Suppose that two 1-links J and K differ by one crossing-change. Then the products of knots, $J \otimes^{\mu}$ (the Hopf link) and*

$K \otimes^{\mu}$ (*the Hopf link*)*, differ by one twist-move, where* $\mu \in \mathbb{N} \cup \{0\}$*. Here,* $A \otimes^{\mu} B$ *means* $A \otimes B ... \otimes B$*, which includes* μ *copies of* B*, where* $\mu \in \mathbb{N} \cup \{0\}$*.*

Note. The fact that the two knots differ means that the two knots are not identical. There are two cases that the two knots are not equivalent and that the two knots are equivalent.

We review the empty knots in §5. It holds that $[2] \otimes [2]$ is the negative Hopf link in S^3 (see [8, 9] and Theorem 5.3.) The above Theorem 2.1 follows from the following Theorem 2.2 by Theorem 5.3.

Theorem 2.2. (Theorem 8.2.) *Take the same* J, K *in Theorem 2.1. Then the* $(2\nu+1)$*-submanifolds* $\subset S^{2\nu+3}$*,* $J \otimes^{\nu}$ (*the empty knot* $[2]$) *and* $K \otimes^{\nu}$ (*the empty knot* $[2]$)*, differ by one twist-move, where* $\nu \in \mathbb{N} \cup \{0\}$*.*

Theorem 2.3. (Theorem 8.3.) Let $m \in \mathbb{N} \cup \{0\}$. Suppose that two $(2m+1)$-dimensional closed oriented submanifolds $\subset S^{2m+3}$, J and K, differ by one twist-move. Then the $(2m+2\nu+1)$-submanifolds $\subset S^{2m+2\nu+3}$, $J \otimes^{\nu}$ (the empty knot $[2]$) and $K \otimes^{\nu}$ (the empty knot $[2]$), differ by one twist-move.

Note. Of course, the $m = 0$ case is Theorem 2.1.

Theorem 2.4. (Theorem 8.6.) *Let* $p \in \mathbb{N}$*. Let* K (*resp.* J) *be* $(2p+5)$*-dimensional smooth submanifold* $\subset S^{2p+7}$*. Suppose that* K *is a spherical knot. Suppose that* K *and* J *differ by a single twist-move and are nonequivalent. Suppose that* K *is equivalent to*

$$\begin{cases} A \otimes^{\frac{p}{2}+1} (\textit{the Hopf link}) \textit{ for a 1-knot } A & \textit{if } p \textit{ is even} \\ A \otimes (\textit{the Hopf link}) \textit{ for a simple 3-knot } A & \textit{if } p = 1 \ (\textit{and hence } 2p+5 = 7) \\ A \otimes^{\frac{p-1}{2}} (\textit{the Hopf link}) \textit{ for a simple 7-knot } A & \textit{if } p \textit{ is odd and } p \neq 1. \end{cases}$$

Then there is a unique equivalence class of simple $(2p+1)$*-knots for* K (*resp.* J) *with the following properties.*

(i) *There is a representative element* K' *of the above equivalence class for* K *such that* K *is equivalent to* $K' \otimes$ (*the Hopf link*).

(ii) *There is a representative element* J' *of the above equivalence class for* J *such that* J *is equivalent to* $J' \otimes$ (*the Hopf link*).

(iii) K' *and* J' *differ by a single twist-move and are nonequivalent.*

Note. Let K be an n-dimensional spherical knot $\subset S^{n+2}$. If $\pi_1(S^{n+2} - K) = \mathbb{Z}$ and if $\pi_i(S^{n+2} - K) = 0 \quad (2 \leqq i < \frac{n}{2}, i \in \mathbb{N})$, then we call K a *simple knot.* See [16].

Note. There are countably infinitely many $(2p+5)$-dimensional spherical knots ($p \in \mathbb{N}$) which are not the product of any $(2p+1)$-knot and the Hopf link by [8, 9].

Note. If $p = 0$, we have a different situation: There are nonequivalent 1-knots K' and J' with the following properties.

(1) K' and J' differ by two crossing-changes not by a crossing-change.

(2) $K'\otimes^\mu$(the Hopf link) and $J'\otimes^\mu$(the Hopf link) differ by a crossing-change and are nonequivalent. (As we stated before, the twist-move on 1-links is the crossing-change on 1-links.)
See Theorem 8.4.

We also prove relations between the pass-move on 1-links and the (p,q)-pass-move on high dimensional knots.

Theorem 2.5. (Theorem 9.1.) *Suppose that two 1-knots J and K differ by one pass-move. Then the $(4\mu+1)$-submanifolds $\subset S^{4\mu+3}$, $J\otimes^\mu$ (the Hopf link) and $K\otimes^\mu$ (the Hopf link), differ by one $(2\mu+1,2\mu+1)$-pass-move $(\mu\in\mathbb{N}\cup\{0\})$.*

Theorem 2.6. (Theorem 9.5.) *Let J, K be simple $(2l+1)$-knots, where $l\in\mathbb{N}$. Suppose that J and K differ by one $(l+1,l+1)$-pass-move. Then the $(2l+4\mu+1)$-submanifolds $\subset S^{2l+4\mu+3}$, $J\otimes^\mu$ (the Hopf link) and $K\otimes^\mu$ (the Hopf link), differ by one $(l+2\mu+1,l+2\mu+1)$-pass-move.*

Theorem 2.7. (Theorem 9.11.) *Let $\mu\in\mathbb{N}$. Let K be a spherical $(4\mu+1)$-knot. Let J be a $(4\mu+1)$-submanifold $\subset S^{4\mu+3}$. Let K and J be $(2\mu+1,2\mu+1)$-pass-move equivalent. Suppose that K is equivalent to $K'\otimes^\mu$ (the Hopf link) for a 1-knot K'. Then there is a 1-knot J' with the following properties.*
(i) *J is equivalent to $J'\otimes^\mu$ (the Hopf link).*
(ii) *K' and J' are pass-move equivalent.*

Theorem 2.8. (Theorem 9.12.) *Let $p\in\mathbb{N}$. Let K and J be $(2p+5)$-dimensional smooth submanifolds $\subset S^{2p+7}$. Suppose that K is a spherical knot. Suppose that K and J differ by a single $(p+3,p+3)$-pass-move and are nonequivalent. Suppose that K is equivalent to*

$$\text{to}\begin{cases} A\otimes^{\frac{p}{2}+1}\,(\textit{the Hopf link})\ \textit{for a 1-knot } A & \textit{if } p \textit{ is even}\\ A\otimes(\textit{the Hopf link})\ \textit{for a simple 3-knot } A & \textit{if } p=1\ (\textit{and hence } 2p+5=7)\\ A\otimes^{\frac{p-1}{2}}\,(\textit{the Hopf link})\ \textit{for a simple 7-knot } A & \textit{if } p \textit{ is odd and } p\neq 1.\end{cases}$$

Then there is a unique equivalence class of simple $(2p+1)$-knots for K (resp. J) with the following properties.
(i) *There is a representative element K' of the above equivalence class for K such that K is equivalent to $K'\otimes$ (the Hopf link).*
(ii) *There is a representative element J' of the above equivalence class for J such that J is equivalent to $J'\otimes$ (the Hopf link).*
(iii) *K' and J' differ by a single $(p+1,p+1)$-pass-move and are nonequivalent.*

Next we discuss a relation between polynomial invariants of 1-links and those of high dimensional knots related by knot products.

Suppose that 1-links K_+, K_-, K_0 differ only in a 3-ball B as shown below.



K_+



K_-



K_0

Then the ordered set (K_+, K_-, K_0) is called a *crossing-change-triple.* We also say that the ordered set (K_+, K_-, K_0) is *related by a crossing-change* in B.

Let $A(K)$ be the Alexander-Conway polynomial of 1-links K. It is well-known that

$$A(K_+) - A(K_-) = (t-1) \cdot A(K_0).$$

Note that there is another kind of setting of the variable. Here, we have the following.

Theorem 2.9. (Theorem 10.1.) *Let K_+, K_-, K_0 be as above. There is a polynomial $\Delta_{2\mu+1}(\ K_* \otimes^\mu$ (the Hopf link) $) \in \mathbb{Q}[t, t^{-1}]$ whose $\mathbb{Q}[t, t^{-1}]$-balanced class is the $(2\mu+1)$-$\mathbb{Q}[t, t^{-1}]$-Alexander polynomial $A_{2\mu+1}(K_* \otimes^\mu$ (the Hopf link)$)$ ($\mu \in \mathbb{N} \cup \{0\}$. $\ * = +, -, 0$.) such that*

$$\begin{aligned} &\Delta_{2\mu+1}(\ K_+ \otimes^\mu (\textit{the Hopf link})\ ) - \Delta_{2\mu+1}(\ K_- \otimes^\mu (\textit{the Hopf link})\ ) \\ &= (t-1) \cdot \Delta_{2\mu+1}(\ K_0 \otimes^\mu (\textit{the Hopf link})\ ). \end{aligned}$$

Note. We review the p-$\mathbb{Q}[t, t^{-1}]$-Alexander polynomial A_p for n-dimensional closed oriented submanifolds $\subset S^{n+2}$ in §6.

The above Theorem 2.9 follows from the following Theorem 2.10 by Theorem 5.3.

Theorem 2.10. (Theorem 10.2.) *Let K_+, K_-, K_0 be as above. There is a polynomial $\Delta_{\nu+1}(K_* \otimes^\nu [2]) \in \mathbb{Q}[t, t^{-1}]$ whose $\mathbb{Q}[t, t^{-1}]$-balanced class is the $(\nu+1)$-$\mathbb{Q}[t, t^{-1}]$-Alexander polynomial $A_{\nu+1}(\ K_* \otimes^\nu [2]\ )$ $\quad(\nu \in \mathbb{N} \cup \{0\}, * = +, -, 0)$ such that*

$$\Delta_{\nu+1}(\ K_+ \otimes^\nu [2]\ ) - \Delta_{\nu+1}(\ K_- \otimes^\nu [2]\ ) = (t + (-1)^{\nu+1}) \cdot \Delta_{\nu+1}(\ K_0 \otimes^\nu [2]\ ),$$

where $[2]$ *denotes the empty knot* $[2]$.

The above Theorem 2.9, 2.10 is connected with the following Theorem 2.11. The 'l =even' case is proved in [23]. In this paper we prove the 'l =odd' case.

Theorem 2.11. (Theorem 10.3.) *Let K_+ be a $(2l+1)$-dimensional spherical knot $\subset S^{2l+3} (l \in \mathbb{N} \cup \{0\})$. Let K_-, K_0 be $(2l+1)$-dimensional submanifolds $\subset S^{2l+3}$. Let (K_+, K_-, K_0) be a twist-move-triple.*

Then there is a polynomial $\Delta_{l+1}(K_*) \in \mathbb{Q}[t, t^{-1}]$ whose $\mathbb{Q}[t, t^{-1}]$-balanced class is the $(l+1)$-$\mathbb{Q}[t, t^{-1}]$-Alexander polynomial $A_{l+1}(K_*)$ $(* = +, -, 0)$ and that

$$\Delta_{l+1}(K_+) - \Delta_{l+1}(K_-) = (t + (-1)^{l+1}) \cdot \Delta_{l+1}(K_0).$$

Note. We define the twist-move-triple in §4. If (K_+, K_-, K_0) is twist-move-triple, then K_+ is obtained from K_-, and K_0 is $[K_+ - (\partial h^{p+1} - \partial B^{n+2})] \cup (h^{p+1} \cap \partial B^{n+2})$. See the following conceptual figure. In the right B, we move $K_0 \cap B$ by isotopy.



This paper is organized as follows.

§1 Introduction
§2 Main results
§3 Local moves on classical links
§4 Local moves on n-knots
§5 Products of knots
§6 Review of the $\mathbb{Q}[t, t^{-1}]$-Alexander polynomials for n-knots and n-dimensional closed oriented submanifolds, etc.
§7 Some results on invariants of n-knots and local moves on n-knots
§8 Theorems on relations between crossing-changes and knot products
§9 Theorems on relations between pass-moves and knot products
§10 Theorems on relations between local move identities of a knot polynomial and knot products
§11 A remark on $\mathbb{Z}[t, t^{-1}]$ case
§12 Proof of Theorems in §8
§13 Proof of Theorems in §9
§14 Proof of Theorems in §10
§15 Discussion and open problems

3. LOCAL MOVES ON CLASSICAL LINKS

Definition 3.1. ([10].) Two 1-links are *pass-move equivalent* if one is obtained from the other by a sequence of pass-moves. See the following figure for an illustrations of the pass-move. Each of four arcs in the 3-ball may belong to different components of the 1-link.



If K and J are pass-move equivalent and if K and K' is equivalent, then we also say that K' and J are pass-move equivalent.

The following proposition is proved in [10].

Theorem 3.2. ([10].) *Let L_1 and L_2 be 1-links. Then L_1 and L_2 are pass-move equivalent if and only if L_1 and L_2 satisfy one of the following conditions (1) and (2).*
(1) Both L_1 and L_2 are proper links, and

$$\mathrm{Arf}(L_1) = \mathrm{Arf}(L_2).$$

(2) Neither L_1 nor L_2 is a proper link, and

$$\mathrm{lk}(K_{1j}, L_1 - K_{1j}) \equiv \mathrm{lk}(K_{2j}, L_2 - K_{2j}) \bmod 2 \quad for \ \ all \ \ j$$

.

4. Local moves on n-knots

We review (p,q)-pass-moves on n-knots ($p, q \in \mathbb{N}$, $\quad p + q = n + 1$) and twist moves on high dimensional knots. [19, 21, 23] defined them. See also [22, 24]. Confirm that, if $(p,q) = (1,1)$, (p,q)-pass-moves are pass-moves on 1-links.

We first define (p,q)-pass-moves on n-knots ($p, q \in \mathbb{N}$, $\quad p + q = n + 1$). Let K_+, K_-, K_0 be n-dimensional closed oriented submanifolds $\subset S^{n+2}$ ($n \in \mathbb{N}$). Let B be an $(n+2)$-ball trivially embedded in S^{n+2}. Suppose that K_+ coincides with K_-, K_0 in $\overline{S^{n+2} - B}$.

Take an $(n+1)$-dimensional p-handle $h^p_*(*=+,-)$ and an $(n+1)$-dimensional $(n+1-p)$-handle h^{n+1-p} in B with the following properties.
(1) $h^p_* \cap \partial B$ is the attaching part of h^p_*. $h^{n+1-p} \cap \partial B$ is the attaching part of h^{n+1-p}.
(2) h^p_* (resp. h^{n+1-p}) is embedded trivially in B.
(3) $h^p_* \cap h^{n+1-p} = \phi$.
(4) The attaching part of h^p_+ coincides with that of h^p_-. The linking number (in B) of
$[h^p_+ \cup (-h^p_-)]$ and
$[h^{n+1-p}$ whose attaching part is fixed in $\partial B]$
is one if an orientation is given.

Let $K_*(*=+,-)$ satisfy that $K_* \cap \mathrm{Int} B = (\partial h^p_* - \partial B) \cup (\partial h^{n+1-p} - \partial B)$.

Let $P = K_+ \cap (S^{n+2} - \mathrm{Int} B)$. Let $Q = h^p_+ \cap \partial B$. Let $R = h^{n+1-p} \cap \partial B$. Let $T = P \cup Q \cup R$. Then T is an n-dimensional oriented closed submanifold $\subset (S^{n+2} - \mathrm{Int} B) \subset S^{n+2}$. Let K_0 be $T \subset S^{n+2}$. Then we say that $(K_+,\ K_-,\ K_0)$ is related by a $(p, n+1-p)$-*pass-move* in B. We also say that $(K_+,\ K_-,\ K_0)$ is a $(p, n+1-p)$-*pass-move-triple*. We say that K_+ and K_- differ by one $(p, n+1-p)$-*pass-move* in B.

If $(K_+,\ K_-,\ K_0)$ is a $(p, n+1-p)$-pass-move-triple, then we also say that $(K_-,\ K_+,\ K_0)$ is a $(p, n+1-p)$-*pass-move-triple*. If K_+ and K_- differ by one $(p, n+1-p)$-pass-move in B, then we also say that K_- and K_+ differ by one $(p, n+1-p)$-*pass-move* in B.

Let (K_+, K_-, K_0) be related by a $(p, n+1-p)$-pass-move in B. Then there is a Seifert hypersurface V_* for K_* $(* = +. - .0)$ with the following properties.
(1) $V_\sharp = V_0 \cup h^p_\sharp \cup h^{n+1-p}$ $(\sharp = +,-)$. $V_\sharp \cap B = h^p_\sharp \cup h^{n+1-p}$.
(2) $V_0 \cap$ Int $B = \phi$. $V_0 \cap \partial B$ is the attaching part of $h^p_\sharp \cup h^{n+1-p}$.
(The idea of the proof is the Thom-Pontrjagin construction.)
Then the ordered set (V_+, V_-, V_0) is called a $(p, n+1-p)$-*pass-move-triple of Seifert hypersurfaces* for (K_+, K_-, K_0). We say that an ordered set (V_+, V_-, V_0) is related by a $(p, n+1-p)$-*pass-move* in B. We say that V_- (resp. V_+) is obtained from V_+ (resp. V_-) by a $(p, n+1-p)$-*pass-move* in B.

Note. When we construct K_- and K_0 from K_+, we make a change only in B and we do not impose any requirement on diffeomorphism type or homeomorphism type of K_-, K_0 other than the change only in B. In this sense, we use the word 'local' in the above definition.

Figure 4.1 is a conceptual figure of a (p, q)-pass-move. Figure 4.2, which consists of the three figures (1) (2) (3), is a conceptual figure of a (p, q)-pass-move-triple.


$B \cap K_+$
$B \cap K_-$
$S^{n-p} \times D^p$
$= \partial h_+^p - \partial B$
$S^{p-1} \times D^{n+1-p}$
$= \partial h^{n+1-p} - \partial B$

Figure 4.1
A $(p, n+1-p)$**-pass-move on an** n**-dimensional submanifold** $\subset S^{n+2}$**. Note** $B = B^{n+2} = D^{n+2} \subset S^{n+2}$**. This figure includes** h_+^p **and** h^{n+1-p}**.**



This cube is $B = D^{n+2} = D^1 \times D^p \times D^{n+1-p}$

$B \cap K_+$

Figure 4.2.(1): **A** $(p, n+1-p)$**-pass-move-triple**



$B \cap K_-$

Figure 4.2.(2): **A** $(p, n+1-p)$**-pass-move-triple**



Figure 4.2.(3): **A** $(p, n+1-p)$**-pass-move-triple**



Figure 4.3.(1): **A (1,2)-pass-move-triple**



Figure 4.3.(2): **A (1,2)-pass-move-triple**



Figure 4.3.(3): **A (1,2)-pass-move-triple**

In Figure 4.3, which consists of the three figures (1) (2) (3), we draw a (1,2)-pass-move-triple. (the $p = 1$ and $n = 2$ case). Since (K_+, K_-, K_0) is related by a $(1, 2)$-pass-move in B, B has the following properties. We regard B as (2-disc)$\times[0, 1] \times \{t| -1 \leqq t \leqq 1\}$.
(i) $K_+ - B$, $K_- - B$, and $K_0 - B$ coincide each other.
(ii) $B \cap K_+$, $B \cap K_-$, $B \cap K_0$ are shown as above.

In the above figures we draw $B_{-0.5} \cap K_*$, $B_0 \cap K_*$, $B_{0.5} \cap K_*$, where B_{t_0}=(2-disc)$\times[0, 1] \times \{t|t = t_0\}$. We suppose that each vector $\overrightarrow{x}$, $\overrightarrow{y}$ in the above figures is a tangent vector of each disc at a point. (Note that we use $\overrightarrow{x}$ (resp. $\overrightarrow{y}$) for different vectors.) The orientation of each disc in the above figures is determined by the each set $\{\overrightarrow{x}, \overrightarrow{y}\}$. Around Figure 4.1 and 4.2 in [21], we wrote more explanation of the figure of $B \cap K_+$ and that of $B \cap K_-$.

In [21] one more local move was discussed, called the 'ribbon-move'. In [21] is proved the following results. Let K and K' be 2-dimensional closed oriented submanifolds $\subset S^4$. The following conditions (1) and (2) are equivalent.
(1) K is (1,2)-pass-move equivalent to K'.
(2) K is ribbon-move equivalent to K'.

Furthermore, if K is obtained from K' by one ribbon-move, then K is obtained from K' by one (1,2)-pass-move.

We next define twist-moves on high dimensional knots. Let K_+, K_-, K_0 be $(2p+1)$-dimensional closed oriented submanifold $\subset S^{2p+3}$ ($p \in \mathbb{N} \cup \{0\}$). Let B be a $(2p+3)$-ball trivially embedded in S^{2p+3}. Suppose that K_+ coincides with K_-, K_0 in $\overline{S^{2p+3} - B}$. Take a single $(2p+2)$-dimensional $(p+1)$-handle h_+ (resp. h_-) embedded in B such that [the handle]$\cap \partial B$ is the attaching part of the handle. Note. [4, 5, 30, 31] etc. imply that the core of h_+ (resp. h_-) is trivially embedded in B under the above condition. Suppose that (h_+-its attaching part)$\cap$(h_--its attaching part)$= \phi$. Suppose that their attaching parts coincide. Thus we can suppose that we can regard $h_+ \cup h_-$ as an oriented submanifold $\subset S^{2p+1}$ if we give the opposite orientation to h_-. Then we can define a $(p+1)$-Seifert matrix for the $(2p+2)$-submanifold $h_+ \cup h_-$. We can suppose that the Seifert matrix is the matrix (1).

Let $K_*(* = +, -)$ satisfy that $K_* \cap \mathrm{Int} B = (\partial h_* - \partial B)$. Note the following. When we define K_+, h_+exists in B and h_- does not exist in B. When we define K_-, h_-exists in B and h_+ does not exist in B. Let $P = K_+ \cap (S^{2p+3} - \mathrm{Int} B)$. Let $Q = h_+ \cap \partial B$. Let $T = P \cup Q$. Then T is an $(2p+1)$-dimensional oriented closed submanifold in $S^{2p+3} - \mathrm{Int} B$. Let K_0 be T in S^{2p+3}. Then we say that an ordered set (K_+, K_-, K_0) is related by a *twist-move.* (K_+, K_-, K_0) is called a *twist-move-triple.* We say that K_+ and K_- differ by one *twist-move* in B. If (K_+, K_-, K_0) is a twist-move-triple, then we also say that (K_-, K_+, K_0) is a *twist-move-triple.* If K_+ and K_- differ by one twist-move in B, we also say that K_- and K_+ differ by one *twist-move* in B.

Let (K_+, K_-, K_0) be related by a twist-move in B. Then there is a Seifert hypersurface V_* for K_* ($* = +. - .0$) with the following properties.
(1) $V_\sharp = V_0 \cup h_\sharp$ ($\sharp = +, -$). $V_\sharp \cap B = h_\sharp$.
(2) $V_0 \cap$ Int $B = \phi$. $V_0 \cap \partial B$ is the attaching part of $h^p_\sharp$.
(The idea of the proof is the Thom-Pontrjagin construction.)

The ordered set (V_+, V_-, V_0) is called a *twist-move-triple of Seifert hypersurfaces* for (K_+, K_-, K_0). We say that V_- (resp. V_+) is obtained from V_+ (resp. V_-) by a *twist-move* in B.

Figure 4.4, which consists of the three figures (1) (2) (3), is a conceptual figure of a twist-move-triple. The upper half of Figure 4.5 is another conceptual figure of a twist-move triple. Compare the upper half of Figure 4.5 and the lower half. If $p = 0$ (hence $n = 2p + 1 = 1$), the left figure in the upper half and that in the lower half are same. That is, if $p = 0$ (hence $n = 2p+1 = 1$), a twist-move-triple is a crossing-change-triple of 1-links. Note that we move $B \cap K_0$ by isotopy in the right B in the upper half of Figure 4.5.



This cube is $D^{2p+3} = B$.

$B \cap K_+$

Figure 4.4.(1): **A twist-move-triple**



Figure 4.4.(2): **A twist-move-triple**



Figure 4.4.(3): **A twist-move-triple**



If $p = 0$



The triple of three ◌ makes a crossing-change-triple of a 1-dimensional link.

Figure 4.5

A twist-move-triple of 1-links is a crossing-change-triple of 1-links.

5. PRODUCTS OF KNOTS

We review the knot product construction as defined in [8, 9].

A *Seifert hypersurface* for an n-dimensional oriented closed submanifold K in S^{n+2} is an $(n+1)$-dimensional oriented connected compact submanifold in S^{n+2} whose boundary is K. Note that there are two cases that K is not connected and that K is connected.

Definition 5.1. ([8, 9].) Let J be an m-dimensional closed oriented submanifold $\subset S^{m+2} = \partial B^{m+3} \subset B^{m+3}$ ($m \in \mathbb{N} \cup \{0\}$). Let K be an n-dimensional closed oriented submanifold $\subset S^{n+2} = \partial B^{n+3} \subset B^{n+3}$ ($n \in \mathbb{N} \cup \{0\}$). Suppose that J or K is a fibered submanifold.

K is called a *fibered submanifold* if K satisfies the following conditions.
(1) $f : S^n - K \to S^1$ is a fiber bundle over S^1.
(2) Let $N(K) = D^2 \times K$ be the tubular neighborhood of K in S^{n+2}. The restriction of $f : S^{n+2} - K \to S^1$ to $N(K)$ is the trivial fibration of $(D^2 \times K - \{0\} \times K) \to S^1$, which is defined by the natural maps $(D^2 \times K - \{0\} \times K) = (D^2 - \{0\}) \times K \to D^2 - \{0\} \to S^1$. Note that each fiber restricted to $S^n -$(the interior of $N(K)$) makes a Seifert hypersurface for K.

We define a *product of the knots* K and J as follows.

Let V be a Seifert hypersurface for J. Note that $J = \partial V \subset V \subset S^{m+2} = \partial B^{m+3} \subset B^{m+3}$. Let W be a Seifert hypersurface for K. Note that $K = \partial W \subset W \subset S^{n+2} = \partial B^{n+3} \subset B^{n+3}$. Push V into B^{m+3}, fixing $\partial V = J$, call this submanifold $\subset B^{m+3}$, V'. Push W into B^{n+3}, fixing $\partial W = K$, call this submanifold $\subset B^{n+3}$, W'. Let $B^2 = \{(a, b) | a \in \mathbb{R}, b \in \mathbb{R}, a^2 + b^2 \leqq 1.\}$. Let $f : B^{m+3} \to B^2$ be a smooth map which satisfies the condition that $f^{-1}((0, 0)) = V'$. Let $g : B^{n+3} \to B^2$ be a smooth map which satisfies the condition that $g^{-1}((0, 0)) = W'$.

Let $M = \{(x, y) \in B^{m+3} \times B^{n+3} | f(x) - g(y) = (0, 0) \in B^2\}$. The $(m+n+3)$-dimensional closed oriented submanifold $\partial M \subset \partial(B^{m+3} \times B^{n+3}) = S^{m+n+5}$ is called *the product of the knots* J and K and is denoted by $J \otimes K$. The product of knots is sometimes called the *knot product*. Strictly speaking, M might have a singularity and hence we must be careful to define the boundary of M. See [8, 9] for detail.

Let $A \otimes^\mu B$ mean $A \otimes B, ..., \otimes B$, which is composed of one copy of A and μ copies of B, where $\mu \in \mathbb{N} \cup \{0\}$. Let $\otimes^\mu B$ mean $B \otimes, ..., \otimes B$, which is composed of μ copies of B, where $\mu \in \mathbb{N} \cup \{0\}$.

Definition 5.2. ([8, 9].) Let $n \in \mathbb{N}$. The *empty knot* $[n]$ is a smooth map $S^1 \to S^1$ such that $\theta \mapsto n\theta$, where $S^1 = \{e^{2\pi i\theta} | \theta \in \mathbb{R}\}$

We regard a Seifert hypersurface of the empty knot $[n]$ as a set of n points $\subset S^1$. We can regard the empty knot $[n]$ as a fibred knot. In [8, 9] is defined a knot product of the empty knot and an n-dimensional closed oriented submanifold $\subset S^{n+2}$.

The (*positive*) *Hopf link* or the (*linking number*) (+1) *Hopf link* is shown as follows.



The *negative Hopf link* or the (*linking number*) (-1) *Hopf link* is shown as follows.



Theorem 5.3. ([8, 9].) *Let* $[n]$ *denote the empty knot of degree* n*. Then we have* $[2] \otimes [2] =$*the negative Hopf link.*

For $\mu \in \mathbb{N}$*,* $\overset{2\mu}{\otimes}\ [2] = \overset{\mu}{\otimes}$ *(the negative Hopf link).*

For any n*-dimensional closed oriented submanifold* $K \subset S^{n+2}$*,*
$K \overset{2\mu}{\otimes}\ [2] = K \overset{\mu}{\otimes}$ *(the negative Hopf link).*

Note. See line(-12) of page 389 and line 18 of page 391 of [9]

In this paper the Hopf link means the negative Hopf link.

6. Review of the $\mathbb{Q}[t, t^{-1}]$-Alexander polynomials for n-knots and n-dimensional closed oriented submanifolds, etc.

We review the $\mathbb{Q}[t, t^{-1}]$-Alexander polynomials for n-knots and n-links and n-dimensional closed oriented submanifolds, Seifert matrices, Alexander matrices, etc. See [1, 14, 15, 16].

Let $K = (K_1, ..., K_\xi)$ be an n-dimensional closed oriented submanifold of S^{n+2} ($n \in \mathbb{N}$). Let each K_i be connected. It is known that any tubular neighborhood of K is diffeomorphic to $K \times D^2$ (see pages 49, 50 of [13]). Let $X = \overline{S^{n+2} - K \times D^2}$. By using the orientation of S^{n+2} and that of K, we can determine an orientation of ∂D^2. Take a homomorphism $\alpha : H_1(X; \mathbb{Z}) \to \mathbb{Z}$ to carry all $[\partial D^2]$ with the orientations to $+1$. Take the infinite cyclic covering $\pi : \widetilde{X} \to X$ associated with α. $\widetilde{X}$ is called the *canonical cyclic* *covering space* of K. We can regard $H_p(\widetilde{X};\mathbb{Z})$ as a $\mathbb{Z}[t,t^{-1}]$-module by using the covering translation $\widetilde{X}\to\widetilde{X}$ defined by α. It is called the $\mathbb{Z}[t,t^{-1}]$-*p-Alexander module.* We can also regard $H_p(\widetilde{X};\mathbb{Q})$ as a $\mathbb{Q}[t,t^{-1}]$-module. It is called the $\mathbb{Q}[t,t^{-1}]$-*p-Alexander module.*

According to module theory, it holds that any $\mathbb{Q}[t,t^{-1}]$-module is congruent to

$$(\mathbb{Q}[t,t^{-1}]/\lambda_1)\oplus\cdots\oplus(\mathbb{Q}[t,t^{-1}]/\lambda_l)\oplus(\oplus^k\mathbb{Q}[t,t^{-1}]),$$

where we have the following:
(1) $\lambda_*\in\mathbb{Q}[t,t^{-1}]$ is not zero,
(2) λ_* is not the $\mathbb{Q}[t,t^{-1}]$-balanced class of 1,
(3) k is the rank of the free part.

Two polynomials $f(t),g(t)\in\mathbb{Q}[t,t^{-1}]$ are said to be $\mathbb{Q}[t,t^{-1}]$-*balanced* if there is an integer n and a nonzero rational number r such that $f(t)=r\cdot t^n\cdot g(t)$.

Let $H_p(\widetilde{X};\mathbb{Q})$ be as above. Then the $\mathbb{Q}[t,t^{-1}]$-*p-Alexander polynomial* is

$$\begin{cases}\text{the } \mathbb{Q}[t,t^{-1}]\text{-balanced class of the product } \lambda_1\cdot\ldots\cdot\lambda_l & \text{if } k=0 \text{ and } H_p(\widetilde{X};\mathbb{Q}) \text{ is nontrivial}\\ 0 & \text{if } k\neq 0\\ 1 & \text{if } H_p(\widetilde{X};\mathbb{Q})\cong 0.\end{cases}$$

Let V be a Seifert hypersurface for the above n-submanifold K. Note that the orientation of V is compatible with that of K. Recall that Seifert hypersurfaces are connected by the definition (see §5). Let $x_1,...,x_\mu$ be p-cycles in V which are basis of $H_p(V;\mathbb{Z})/\text{Tor}$. Let $y_1,...,y_\nu$ be $(n+1-p)$-cycles in V which are basis of $H_p(V;\mathbb{Z})/\text{Tor}$. Push y_i to the positive direction of the normal bundle of V. Call it y_i^+. Push y_i to the negative direction of the normal bundle of V. Call it y_i^-. A $(p,n+1-p)$-*(positive) Seifert matrix* for the above submanifold K associated with V represented by an 'ordered set of basis', $\{x_1,...,x_\mu\}$, and an 'ordered set of basis', $\{y_1,...,y_\nu\}$, is a $(\mu\times\nu)$-matrix $S=(s_{ij})=(\text{lk}(x_i,y_j^+))$. We sometimes abbreviate $(p,n+1-p)$-Seifert matrix to p-Seifert matrix if we know what n is. We sometimes let $S_p(K)$ denote a positive p-Seifert matrix for a closed oriented submanifold K and V and $\{x_i\}$ and $\{y_j\}$ if we know what V and $\{x_i\}$ and $\{y_j\}$ are.

A $(p,n+1-p)$-*negative Seifert matrix* for the above submanifold K associated with V represented by an 'ordered set of basis', $\{x_1,...,x_\mu\}$, and an 'ordered set of basis', $\{y_1,...,y_\nu\}$, is a matrix $N=(n_{ij})=(\text{lk}(x_i,y_j^-))$. We sometimes let $N_p(V)$ denote a negative p-Seifert matrix for a closed oriented submanifold K and V and $\{x_i\}$ and $\{y_j\}$ if we know what K and $\{x_i\}$ and $\{y_j\}$ are. Let S_p and N_p be as above. Then we have the following. S_p-N_p represents the map $\{H_p(V;\mathbb{Z})/\text{Tor}\}\times\{H_{n+1-p}(V;\mathbb{Z})/\text{Tor}\}\to\mathbb{Z}$, which is defined by the intersection product. We call $t\cdot S_p-N_p$ the *p-Alexander matrix* for K associated with V represented by an 'ordered set of basis', $\{x_1,...,x_\mu\}$, and an 'ordered set of basis', $\{y_1,...,y_\nu\}$.

Note that we sometimes define it to be $S_p - t \cdot N_p$. The difference of both is only setting the variables because we mainly discuss $\mathbb{Q}[t, t^{-1}]$-balanced-classes as follows. All we have to do is to change t with t^{-1}.

We have the following.

Proposition 6.1. *Let K be an n-dimensional oriented closed submanifold $\subset S^{n+2}$.*

Let S_p be a $(p, n+1-p)$-positive Seifert matrix for K associated with V represented by an 'ordered set of basis', $\{x_1, ..., x_\mu\}$, and an 'ordered set of basis', $\{y_1, ..., y_\nu\}$.

Let N_p be a $(p, n+1-p)$-negative Seifert matrix for K associated with V represented by an 'ordered set of basis', $\{x_1, ..., x_\mu\}$, and an 'ordered set of basis', $\{y_1, ..., y_\nu\}$.

Suppose $\mu = \nu$. Suppose that the linear map defined by a $(p-1)$-Alexander matrix is injective.

Then the p-$\mathbb{Q}[t, t^{-1}]$-Alexander polynomial is the $\mathbb{Q}[t, t^{-1}]$-balanced class of 'the determinant of p-Alexander matrix,' $\det\,(t \cdot S_p - N_p)$.

Note. Of course, $\mu \neq \nu$ in general.

Proof. Take the above $X = \overline{S^{n+2} - K \times D^2}$, $\widetilde{X}$, V. Let $V \times [-1, 1]$ be the tubular neighborhood of V in X. Let $Y = X - V$. Consider the Meyer-Vietoris exact sequence:
$H_\natural(\amalg_{-\infty}^{\infty} V \times [-1, 1]; \mathbb{Q}) \to H_\natural(\amalg_{-\infty}^{\infty} Y; \mathbb{Q}) \to H_\natural(\widetilde{X}; \mathbb{Q})$,
where $\amalg_{-\infty}^{\infty} V \times [-1, 1]$ is the lift of $V \times [-1, 1]$, and where $\amalg_{-\infty}^{\infty} Y$ is the lift of Y. This completes the proof. $\square$

We have the following.

Let N_p be a $(p, n+1-p)$-negative Seifert matrix for K associated with V represented by an 'ordered set of basis', $\{x_1, ..., x_\mu\}$, and an 'ordered set of basis', $\{y_1, ..., y_\nu\}$. Let S_{n+1-p} be a $(n+1-p, p)$-positive Seifert matrix for K associated with V represented by an 'ordered set of basis', $\{y_1, ..., y_\nu\}$, and an 'ordered set of basis', $\{x_1, ..., x_\mu\}$. By the definition of x_i^+ and y_i^-, $\mathrm{lk}(y_i, x_j^+) = \mathrm{lk}(y_i^-, x_j)$. By page 541 of [14], $\mathrm{lk}(y_i^-, x_j)$ $= (-1)^{p(n+1-p)+1}\mathrm{lk}(x_j, y_i^-)$. Hence $N_p = (-1)^{p \cdot n+1} S_{n+1-p}$ (note that $p(1-p)$ is an even number).

Proposition 6.2. *Let K be a $(2m+1)$-dimensional closed oriented submanifold $\subset S^{2m+3}$. Let S be an $(m+1, m+1)$-Seifert matrix. We have $S = (-1)^m \cdot {}^t S$.*

The *signature* $\sigma(K)$ for a $(2p+1)$-dimensional closed oriented submanifold K $\subset S^{2p+3} (p \in \mathbb{N} \cup \{0\})$ is the signature of the matrix $S_{p+1}(K) + {}^t S_{p+1}(K)$. Therefore, we have the following.

Claim 6.3. *Let K be a $(4k+3)$-dimensional closed oriented submanifold $\subset S^{4k+5}$ $(k$ $\in \mathbb{N} \cup \{0\})$. Then the signature of K coincides with the signature of $\hat{V}$, where $\hat{V}$ is the closed oriented manifold which we obtain by attaching a $(4k+4)$-dimensional 0-handles to ∂V.*

Let K be a $(4k+1)$-dimensional spherical knot $(4k+1 \geq 1. \quad k \in \mathbb{N} \cup \{0\})$. We regard naturally $(H_{2k+1}(V;\mathbb{Z})/\mathrm{Tor}) \otimes \mathbb{Z}_2$ as a subgroup of $H_{2k+1}(V;\mathbb{Z}_2)$. Then we can take basis $x_1, ..., x_\nu, y_1, ..., y_\nu$ of $(H_{2k+1}(V;\mathbb{Z})/\mathrm{Tor}) \otimes \mathbb{Z}_2$ such that $x_i \cdot x_j = 0$, $y_i \cdot y_j = 0$, $x_i \cdot y_j = \delta_{ij}$ for any pair (i,j), where $\cdot$ denotes the $\mathbb{Z}_2$-intersection product. The *Arf invariant* of K is mod 2 $\Sigma_{i=1}^{\nu} \mathrm{lk}(x_i, x_i^+) \cdot \mathrm{lk}(y_i, y_i^+)$.

Let $L = (L_1, ..., L_\mu)$ be a $(4k+1)$-link $(4k+1 \geq 1. \quad k \in \mathbb{N} \cup \{0\}. \quad \mu \in \mathbb{N} - \{0\}.)$. We define the Arf invariant of L. There are two cases.

(1) Let $4k+1 \geq 5$. The Arf invariant of L is defined in the same manner as the knot case.

(2) Let $4k+1 = 1$. See Appendix of [13] and Note right above Note 1.2.1 of [20].

7. Some results on invariants of n-knots and local moves on n-knots

Theorem 7.1. ([23].) *Let K_+, K_- be spherical n-knots $\subset S^{n+2}$ $(n \in \mathbb{N})$. Let K_0 be an n-submanifold $\subset S^{n+2}$. Suppose that $(K_+,\ K_-,\ K_0)$ is related by a $(p, n+1-p)$-pass-move. Let $p \neq n+1-p$. Then there is a polynomial $\Delta_p(K_*) \in \mathbb{Q}[t,t^{-1}]$ whose $\mathbb{Q}[t,t^{-1}]$-balanced class is the p-$\mathbb{Q}[t,t^{-1}]$-Alexander polynomial $A_p(K_*)$ for the submanifold K_* $(* = +, -, 0)$ such that*

$$\Delta_p(K_+) - \Delta_p(K_-) = (t-1) \cdot \Delta_p(K_0).$$

Theorem 7.2. ([23].) *Let K_+, K_- be spherical $(4k+1)$-knots $(4k+1 \in \mathbb{N}, \quad k \in \mathbb{N} \cup \{0\})$. Let K_0 be a $(4k+1)$-submanifold $\subset S^{4k+3}$. Suppose that $(K_+,\ K_-,\ K_0)$ is related by a twist-move.*

Then there is a polynomial $\Delta_{2k+1}(K_*) \in \mathbb{Q}[t,t^{-1}]$ whose $\mathbb{Q}[t,t^{-1}]$-balanced class is the $(2k+1)$-$\mathbb{Q}[t,t^{-1}]$-Alexander polynomial $A_{2k+1}(K_*)$ $(* = +, -, 0)$ such that

$$\Delta_{2k+1}(K_+) - \Delta_{2k+1}(K_-) = (t-1) \cdot \Delta_{2k+1}(K_0).$$

Note. For $(4k+3)$-knots $(4k+3 \in \mathbb{N}, \quad k \in \mathbb{N} \cup \{0\})$, we can define the twist-move. However, note the following: Suppose K_+ is made into K_- by the twist-move. Then K_- is not spherical in general. Because: there is an example V_* for K_* as follows; Framed link representation of V_+ is the Hopf link such that the framing of one component is zero and that that of the other is two. Framed link representation of V_- is the Hopf link such that the framing of each component is two.

We prove the following Theorem 2.11 (Theorem 10.3). The 'l =even' case is proved in [23], which is the above Theorem 7.2. In §14 of this paper we prove the 'l =odd' case.

Theorem 2.11. (Theorem 10.3.) *Let K_+ be a $(2l+1)$-dimensional spherical knot $\subset S^{2l+3} (l \in \mathbb{N} \cup \{0\})$. Let K_-, K_0 be $(2l+1)$-dimensional submanifolds $\subset S^{2l+3}$. Let (K_+, K_-, K_0) be a twist-move-triple. Then there is a polynomial $\Delta_{l+1}(K_*) \in \mathbb{Q}[t,t^{-1}]$*

whose $\mathbb{Q}[t,t^{-1}]$*-balanced class is the* $(l+1)$*-*$\mathbb{Q}[t,t^{-1}]$*-Alexander polynomial* $A_{l+1}(K_*)$ $(*=+,-,0)$ *and that*

$$\Delta_{l+1}(K_+) - \Delta_{l+1}(K_-) = (t + (-1)^{l+1}) \cdot \Delta_{l+1}(K_0).$$

Theorem 7.3. ([23].) *Let* K_+*,* K_-*,* K_0 *be as in Theorem 7.2. Let* bP_{4k+2} *be the* bP*-subgroup* $\subset \Theta^{4k+1}$*. Suppose* bP_{4k+2} *is not congruent to the trivial group. Then we have*

$$\mathrm{Arf}K_+ - \mathrm{Arf}K_- = \{|bP_{4k+2} \cap I(K_0)| + 1\} \ \mathrm{mod}2,$$

where $I(K_0)$ *is the inertia group of an oriented smooth manifold which is orientation preserving diffeomorphic to* K_0 *and the symbol* $| \quad |$ *denotes the order of a group.*

Note. See §6 and [12, 13, 14] for the Arf invariant. See [12] for the homotopy sphere group $\Theta^\star$ and the bP-subgroup $\subset \Theta^{4k+1}$. See [2, 11] for the inertia group.

We state a problem.

Problem 7.4. Let K_+ be an n-dimensional spherical knot. Suppose that (K_+, K_-, K_0) is a twist-move-triple (resp. $(p, n+1-p)$-pass-move-triple, where $p \neq n+1-p$). Let $\alpha(K)$ be an invariant of K as a submanifold and be a $\mathbb{Q}[t,t^{-1}]$-balanced class. Suppose that there are $f_+, f_-, f_0 \in \mathbb{Q}[t,t^{-1}]$ such that the $\mathbb{Q}[t,t^{-1}]$-balanced class of f_* is $\alpha(K_*)$ $(*=+,-,0)$ and that

$$\begin{cases} f_+ - f_- = (t-1)\cdot f_0 & \text{in the other cases than the following} \\ f_+ - f_- = (t+1)\cdot f_0 & \text{if } n = 4k+3,\ k \in \mathbb{N}\cup\{0\} \text{ and if we consider the twist-move.} \end{cases}$$

Then is $\alpha(K)$ the $\mathbb{Q}[t,t^{-1}]$-Alexander polynomial or what is determined by the $\mathbb{Q}[t,t^{-1}]$-Alexander polynomial?

Note. It is well-known that the Alexander-Conway polynomial (resp. the Jones polynomial) of 1-links is essentially characterized by the well-known local move formula and the fact that it is trivial for the trivial knot.

8. THEOREMS ON RELATIONS BETWEEN CROSSING-CHANGES AND KNOT PRODUCTS

Theorem 8.1. *Suppose that two 1-links* J *and* K *differ by one crossing-change. Then the 3-submanifolds* $\subset S^5$*,* $J \otimes [2]$ *and* $K \otimes [2]$*, differ by one twist-move, where* $[2]$ *denotes the empty knot* $[2]$*.*

Theorem 8.2. (Theorem 2.2.) *Take the same* J, K *in Theorem 8.1. Then the* $(2\nu+1)$*-submanifolds* $\subset S^{2\nu+3}$*,* $J \otimes^\nu [2]$ *and* $K \otimes^\nu [2]$*, differ by one twist-move, where* $\nu \in \mathbb{N}\cup\{0\}$*, and, where* $[2]$ *denotes the empty knot* $[2]$*.*

Note. It is true even in the $\nu = 0$ case. Of course the $\nu = 1$ case is Theorem 8.1.

Theorem 8.3. (Theorem 2.3.) *Let* $m \in \mathbb{N} \cup \{0\}$*. Suppose that two* $(2m+1)$*-dimensional closed oriented submanifolds* $\subset S^{2m+3}$*,* J *and* K*, differ by one twist-move. Then the* $(2m+2\nu+1)$*-submanifolds* $\subset S^{2m+2\nu+3}$*,* $J \otimes^\nu [2]$ *and* $K \otimes^\nu [2]$*, differ by one twist-move, where* $[2]$ *denotes the empty knot* $[2]$*.*

Note. Of course, the $m = 0$ case is Theorem 8.2.

Theorem 8.4. (1) *There is a nontrivial 1-knot* K *which is obtained from the trivial knot by a single crossing-change with the following property. The* $(2\nu+1)$*-submanifold* $\subset S^{2\nu+3}$*,* $K \otimes^\nu [2]$*, is equivalent to the trivial* $(2\nu+1)$*-knot, where* $\nu \in \mathbb{N}, \quad \nu \geqq 2$*, and, where* $[2]$ *denotes the empty knot* $[2]$*.*

(2) *There is a nontrivial 1-knot* P *which is obtained from the trivial knot by two crossing-changes not by a single crossing-change with the following property. The* $(2\nu+1)$*-submanifold* $\subset S^{2\nu+3}$*,* $P \otimes^\nu [2]$*, is equivalent to the trivial* $(2\nu+1)$*-knot, where* $\nu \in \mathbb{N}, \quad \nu \geqq 2$*.*

(3) *There are nontrivial 1-knots* P *and* Q *with the following properties.*
(i) P *and* Q *differ by a single crossing-change and are nonequivalent.*
(ii) *Let* $\nu \in \mathbb{N}$ *and* $\nu \geqq 2$*. The* $(2\nu+1)$*-submanifolds* $\subset S^{2\nu+3}$*,* $P \otimes^\nu [2]$ *and* $Q \otimes^\nu [2]$ *are equivalent spherical knots.*

(4) *There is a nontrivial 1-knot* P *with the following properties.*
(i) P *is obtained from the trivial 1-knot by two crossing-changes not by a single crossing-change.*
(ii) *Let* $\mu \in \mathbb{N}$*. The* $(4\mu+1)$*-submanifold* $\subset S^{4\mu+3}$*,* $P \otimes^\mu$ (*the Hopf link*) *is obtained from the trivial knot by a twist-move and is a nontrivial knot.*

(5) *There are nontrivial 1-knots* P *and* Q *with the following properties.*
(i) P *and* Q *differ by a single crossing-change and are nonequivalent.*
(ii) *Let* $\mu \in \mathbb{N}$*. The* $(4\mu+1)$*-submanifolds* $\subset S^{4\mu+3}$*,* $P \otimes^\mu$ (*the Hopf link*) *and* $Q \otimes^\mu$ (*the Hopf link*) *are equivalent spherical knots and nontrivial knots.*

Note. Recall that, if a 1-knot K is obtained from a nonequivalent 1-knot J by a single crossing-change, then the 1-knot K is obtained from the nonequivalent knot J by a single twist-move. Thus we can say that the 'twist-move-unknotting-number' changes by a knot product. A 'non-twist-move-equivalent pair' is changed into a 'twist-move-equivalent pair' by knot product.

Note 8.5. Compare the above Theorem 8.4 with the following $(*)$.
$(*)$ There is a nontrivial 1-knot K which is obtained from the trivial knot by one crossing-change with the following properties:
(1) $K \otimes [2]$ is not a spherical knot. Hence it is a nontrivial knot.

(2) $K \otimes^{\nu} [2]$ $(\nu \geqq 2, \nu \in \mathbb{N})$ is a spherical knot and the trivial knot.
We prove this in §12.

Furthermore, we prove the following.

Theorem 8.6. (Theorem 2.4) *Let $p \in \mathbb{N}$. Let K (resp. J) be a $(2p+5)$-dimensional smooth submanifold $\subset S^{2p+7}$. Suppose that K is a spherical knot. Suppose that K and J differ by a single twist-move and are nonequivalent. Suppose that K is equivalent to*

$$\begin{cases} A \otimes^{\frac{p}{2}+1} (\textit{the Hopf link}) \textit{ for a 1-knot } A & \textit{if } p \textit{ is even} \\ A \otimes (\textit{the Hopf link}) \textit{ for a simple 3-knot } A & \textit{if } p = 1 \ (\textit{and hence } 2p+5 = 7) \\ A \otimes^{\frac{p-1}{2}} (\textit{the Hopf link}) \textit{ for a simple 7-knot } A & \textit{if } p \textit{ is odd and } p \neq 1. \end{cases}$$

Then there is a unique equivalence class of simple $(2p+1)$-knots for K (resp. J) with the following properties.
(i) *There is a representative element K' of the above equivalence class for K such that K is equivalent to $K' \otimes$ (the Hopf link).*
(ii) *There is a representative element J' of the above equivalence class for J such that J is equivalent to $J' \otimes$ (the Hopf link).*
(iii) *K' and J' differ by a single twist-move and are nonequivalent.*

Note. If $p = 0$ in Theorem 8.6, we have a different result. See Theorem 8.4.

9. Theorems on relations between pass-moves and knot products

Theorem 9.1. (Theorem 2.5.) *Suppose that two 1-knots J and K differ by one pass-move. Then the $(4\mu+1)$-submanifolds $\subset S^{4\mu+3}$, $J \otimes^{\mu}$ (the Hopf link) and $K \otimes^{\mu}$ (the Hopf link), differ by one $(2\mu+1, 2\mu+1)$-pass-move $(\mu \in \mathbb{N} \cup \{0\})$.*

Note 9.2. Recall Theorem 5.3. For any n-dimensional closed oriented submanifold $A \subset S^{n+2}$, $A \overset{\mu}{\otimes}$ (the negative Hopf link) $= A \overset{2\mu}{\otimes} [2]$, where $n \in \mathbb{N}$ and $\mu \in \mathbb{N} \cup \{0\}$.

Compare the above Theorem 9.1 and Note 9.2 with the following Theorem 9.3.

Theorem 9.3. *Take the same J, K in Theorem 9.1. Then the $(4\mu+3)$-submanifolds $\subset S^{4\mu+5}$ $(\mu \in \mathbb{N} \cup \{0\})$, $J \overset{(2\mu+1)}{\otimes} [2]$ and $K \overset{(2\mu+1)}{\otimes} [2]$, are not homeomorphic in general and, therefore, are NOT $(2\mu+2, 2\mu+2)$-pass-move equivalent in general.*

Problem 9.4. In the above Theorem 9.3, of course, if J and K are trivial knots, then the above two $(4\mu+3)$-submanifolds are pass-move equivalent. What kind of pair, J and K, in Theorem 9.3 satisfies the condition that the above two $(4\mu+3)$-submanifolds are $(2\mu+2, 2\mu+2)$-pass-move equivalent?

The 'ν =odd' case of Theorem 9.8 and Note 9.10 give partial solutions to Problem 9.4.

Theorem 9.5. *Let J, K be simple $(2l+1)$-knots, where $l \in \mathbb{N}$. Suppose that J and K differ by one $(l+1, l+1)$-pass-move. Then the $(2l+4\mu+1)$-submanifolds $\subset S^{2l+4\mu+3}$, $J \otimes^{\mu}$(the Hopf link) and $K \otimes^{\mu}$(the Hopf link), differ by one $(l+2\mu+1, l+2\mu+1)$-pass-move.*

Problem 9.6. If we do NOT suppose that J, K are simple knots in Theorem 9.5, do the above $(2l+4\mu+1)$-submanifolds differ by one pass-move? Or, are they pass-move-equivalent?

The above problem is really a problem of high dimensional knots. The following one is also such a problem.

Problem 9.7. (A generalization of Problem 9.6.) (1) Suppose that spherical n-knots (resp. n-dimensional closed oriented submanifolds $\subset S^{n+2}$) J and K differ by one $(p, n+1-p)$-pass-move, where $n \in \mathbb{N}$ and $p \in \mathbb{N}$. Then do the $(n+4\mu+1)$-submanifolds $\subset S^{n+4\mu+3}$, $J \otimes^{\mu}$ (the Hopf link) and $K \otimes^{\mu}$ (the Hopf link), differ by one pass-move? Or, are they pass-move equivalent?
(2) How about the case where J and K are even dimensional simple $2m$-knots and where $p = m$? Here, $m \in \mathbb{N}$.
(3) Of course, there is a problem in the case of the product with odd times copies of the empty knot [2]. (Note Theorem 9.3 and its Proof.)

Theorem 9.8. *There is a nontrivial 1-knot K which is obtained from the trivial knot by one pass-move with the following property: $K \otimes^{\nu}$ [2] is the trivial $(2\nu+1)$-knot, where $\nu \geqq 2, \quad \nu \in \mathbb{N}$.*

Note 9.9. By the above Theorem 9.8, we have the following. Let T be the trivial 1-knot. The two 1-knots, K and T, differ by a single pass-move and are nonequivalent. However, the $(2\nu+1)$-dimensional spherical knot, $K \otimes^{\nu}$ [2], is equivalent to the $(2\nu+1)$-dimensional trivial knot, $T \otimes^{\nu}$ [2]. That is, they differ by ZERO times of pass-moves, where $\nu \geqq 2$ and $\nu \in \mathbb{N}$. Thus we can say that the 'pass-move-unknotting-number' changes by knot products.

Note 9.10. Compare the above Theorem 9.8 with the following $(*)$.
$(*)$ There is a nontrivial 1-knot K which is obtained from the trivial knot by one pass-move with the following properties:
(1) $K \otimes$ [2], is not a spherical knot. Hence it is a nontrivial knot.
(2) $K \otimes^{\nu}$ [2] ($\nu \geqq 2, \nu \in \mathbb{N}$) is a spherical knot and the trivial knot.
We prove this in §13.

Furthermore, we prove the following.

Theorem 9.11. (Theorem 2.7.) *Let $\mu \in \mathbb{N}$. Let K be a spherical $(4\mu+1)$-knot. Let J be a $(4\mu+1)$-submanifold $\subset S^{4\mu+3}$. Let K and J be $(2\mu+1, 2\mu+1)$-pass-move equivalent.*

Suppose that K is equivalent to $K' \otimes^\mu$ (the Hopf link) for a 1-knot K'. Then there is a 1-knot J' with the following properties.
(i) *J is equivalent to $J' \otimes^\mu$ (the Hopf link).*
(ii) *K' and J' are pass-move equivalent.*

Note. If there is such a 1-knot K', there are countably infinitely many nonequivalent 1-knots P (resp. Q) such that $P \otimes^\mu$ (the Hopf link) (resp. $Q \otimes^\mu$ (the Hopf link)) is equivalent to K (resp. J). We prove this in §13.

Theorem 9.12. (Theorem 2.8.) *Let $p \in \mathbb{N}$. Let K and J be $(2p+5)$-dimensional smooth submanifolds $\subset S^{2p+7}$. Suppose that K is a spherical knot. Suppose that K and J differ by a single $(p+3, p+3)$-pass-move and are nonequivalent. Suppose that K is equivalent to*

$$\begin{cases} A \otimes^{\frac{p}{2}+1} \text{(the Hopf link) for a 1-knot } A & \text{if } p \text{ is even} \\ A \otimes \text{(the Hopf link) for a simple 3-knot } A & \text{if } p = 1 \ (\text{and } 2p+5 = 7) \\ A \otimes^{\frac{p-1}{2}} \text{(the Hopf link) for a simple 7-knot } A & \text{if } p \text{ is odd and } p \neq 1. \end{cases}$$

Then there is a unique equivalence class of simple $(2p+1)$-knots for K (resp. J) with the following properties.
(i) *There is a representative element K' of the above equivalence class for K such that K is equivalent to $K' \otimes$ (the Hopf link).*
(ii) *There is a representative element J' of the above equivalence class for J such that J is equivalent to $J' \otimes$ (the Hopf link).*
(iii) *K' and J' differ by a single $(p+1, p+1)$-pass-move and are nonequivalent.*

Let P be the 5-twist spun knot of the trefoil knot. Note that P is a 2-knot. Note that Proposition 4.3 of [21] and the last line of §7 in page 684 of [21] imply the following.
(1) P is NOT ribbon move equivalent to T.
(2) P is NOT (1,2)-pass-move equivalent to T.

Theorem 9.13. *Let T be the trivial 2-knot. Let P be as above. Although P is NOT (1,2)-pass-move equivalent to T, we have the following. Let $\nu \geqq 2$ and $\nu \in \mathbb{N}$. The $(2\nu+2)$-submanifold, $P \otimes^\nu [2]$, is equivalent to the trivial $(2\nu+2)$-knot, $T \otimes^\nu [2]$, and therefore, is $(\nu+1, \nu+2)$-pass-move-equivalent to the trivial knot.*

Note. Thus we can say that a knot product makes a 'non-pass-move-equivalent pair' into a 'pass-move-equivalent pair'.

10. THEOREMS ON RELATIONS BETWEEN LOCAL MOVE IDENTITIES OF A KNOT POLYNOMIAL AND KNOT PRODUCTS

Let (K_+, K_-, K_0) be a crossing-change-triple of 1-links. (See crossing-change-triple for §1.) Let $A(K)$ be the Alexander-Conway polynomial of 1-links K. It is well-known that

$$A(K_+) - A(K_-) = (t-1) \cdot A(K_0).$$

Here, we have the following Theorems.

Theorem 10.1. (Theorem 2.9.) *Let* K_+, K_-, K_0 *be as above. There is a polynomial* $\Delta_{2\mu+1}(\ K_* \otimes^\mu$ *(the Hopf link)* $) \in \mathbb{Q}[t, t^{-1}]$ *whose* $\mathbb{Q}[t, t^{-1}]$*-balanced class is the* $(2\mu + 1)$*-*$\mathbb{Q}[t, t^{-1}]$*-Alexander polynomial* $A_{2\mu+1}(K_* \otimes^\mu$ *(the Hopf link)*$)$ $(\mu \in \mathbb{N} \cup \{0\}. \quad * = +, -, 0.)$ *such that*

$$\begin{aligned} &\Delta_{2\mu+1}(\ K_+ \otimes^\mu \text{(the Hopf link)}\ ) - \Delta_{2\mu+1}(\ K_- \otimes^\mu \text{(the Hopf link)}\ ) \\ &= (t-1) \cdot \Delta_{2\mu+1}(\ K_0 \otimes^\mu \text{(the Hopf link)}\ ). \end{aligned}$$

Note. After taking knot product, $A(\quad)$ is changed into $A_{2\mu+1}(\quad)$.

By Theorem 5.3, Theorem 10.1 follows from Theorem 10.2.

Theorem 10.2. (Theorem 2.10.) *Let* K_+, K_-, K_0 *be as above. There is a polynomial* $\Delta_{\nu+1}(K_* \otimes^\nu [2]) \in \mathbb{Q}[t, t^{-1}]$ *whose* $\mathbb{Q}[t, t^{-1}]$*-balanced class is the* $(\nu+1)$*-*$\mathbb{Q}[t, t^{-1}]$*-Alexander polynomial* $A_{\nu+1}(\ K_* \otimes^\nu [2]\ )$ $(\nu \in \mathbb{N} \cup \{0\}, * = +, -, 0)$ *such that*

$$\Delta_{\nu+1}(\ K_+ \otimes^\nu [2]\ ) - \Delta_{\nu+1}(\ K_- \otimes^\nu [2]\ ) = (t + (-1)^{\nu+1}) \cdot \Delta_{\nu+1}(\ K_0 \otimes^\nu [2]\ ),$$

where $[2]$ *denotes the empty knot* $[2]$.

Note. If ν is odd, then $K_+ \otimes^\nu [2]$ is not homeomorphic to $K_- \otimes^\nu [2]$ in general. However, the above theorem is true.

The following Theorem 10.3 is much connected with Theorem 10.4. The 'l =even' case is proved in [23]. In this paper we prove the 'l =odd' case.

Theorem 10.3. (Theorem 2.11.) *Let* K_+ *be a* $(2l + 1)$*-dimensional spherical knot* $\subset S^{2l+3}(l \in \mathbb{N} \cup \{0\})$. *Let* K_-, K_0 *be* $(2l + 1)$*-dimensional submanifolds* $\subset S^{2l+3}$. *Let* (K_+, K_-, K_0) *be a twist-move-triple.*

Then there is a polynomial $\Delta_{l+1}(K_*) \in \mathbb{Q}[t, t^{-1}]$ *whose* $\mathbb{Q}[t, t^{-1}]$*-balanced class is the* $(l + 1)$*-*$\mathbb{Q}[t, t^{-1}]$*-Alexander polynomial* $A_{l+1}(K_*)$ $(* = +, -, 0)$ *and that*

$$\Delta_{l+1}(K_+) - \Delta_{l+1}(K_-) = (t + (-1)^{l+1}) \cdot \Delta_{l+1}(K_0).$$

Theorem 10.4. *Let* K_+, K_-, K_0 *be as in Theorem 10.3. There is a polynomial* $\Delta_{l+1+\nu}(\ K_* \otimes^\nu [2]\ ) \in \mathbb{Q}[t, t^{-1}]$ *whose* $\mathbb{Q}[t, t^{-1}]$*-balanced class is the* $(l + 1 + \nu)$*-*$\mathbb{Q}[t, t^{-1}]$*-Alexander polynomial* $A_{l+1+\nu}(\ K_* \otimes^\nu [2]\ )$ $(\nu \in \mathbb{N} \cup \{0\}. \quad * = +, -, 0.)$ *such that*

$$\Delta_{l+1+\nu}(\ K_+ \otimes^\nu [2]\ ) - \Delta_{l+1+\nu}(\ K_- \otimes^\nu [2]\ ) = (t + (-1)^{l+1+\nu}) \cdot \Delta_{l+1+\nu}(\ K_0 \otimes^\nu [2]\ ),$$

where $[2]$ *denotes the empty knot* $[2]$.

Theorem 10.5. *Let* K_+ *be a* $(4l + 1)$*-dimensional spherical knot* $(l \in \mathbb{N} \cup \{0\})$. *Let* (K_+, K_-, K_0) *be a twist-move-triple. Let* $bP_{4l+2+4\mu}$ *be the* bP*-subgroup* $\subset \Theta^{4l+1+4\mu}$. *Suppose* $bP_{4l+2+4\mu}$ *is not congruent to the trivial group. Then we have*

$$\begin{aligned}&\mathrm{Arf}(\ K_+ \otimes^\mu (\textit{the Hopf link})\ ) - \mathrm{Arf}(\ K_- \otimes^\mu (\textit{the Hopf link})\ )\\ &= \{|bP_{4k+2} \cap I(\ K_0 \otimes^\mu (\textit{the Hopf link})\ )| + 1\} \bmod 2,\end{aligned}$$

where $I(\ K_0 \otimes^\mu (\textit{the Hopf link})\ )$ *is the inertia group of an oriented smooth manifold which is orientation preserving diffeomorphic to* $K_0 \otimes^\mu$ *(the Hopf link) and the symbol* $|\quad|$ *denotes the order of a group.*

Let K_+ and K_- be n-dimensional spherical knots $\subset S^{n+2}$. Let K_0 be an n-submanifold $\subset S^{n+2}$. Let (K_+, K_-, K_0) be related by a (p, q)-pass-move in B^{n+2}. Let $p + q = n + 1$. Let $p \neq q$. Recall that we have the following by [23]. (It is quoted in Theorem 7.1 in this paper.)

There is a polynomial $\Delta_p(K_*) \in \mathbb{Q}[t, t^{-1}]$ whose $\mathbb{Q}[t, t^{-1}]$-balanced class is the p-$\mathbb{Q}[t, t^{-1}]$-Alexander polynomial $A_p(K_*)$ for the submanifold K_* $(* = +, -, 0)$ such that

$$\Delta_p(K_+) - \Delta_p(K_-) = (t - 1) \cdot \Delta_p(K_0).$$

We have the following.

Theorem 10.6. *Let* K_+, K_-, K_0 *be as above. There is a polynomial* $\Delta_{p+\nu}(\ K_* \otimes^\nu [2]\ ) \in \mathbb{Q}[t, t^{-1}]$ *whose* $\mathbb{Q}[t, t^{-1}]$*-balanced class is the* $(p + \nu)$*-*$\mathbb{Q}[t, t^{-1}]$*-Alexander polynomial* $A_{p+\nu}(\ K_* \otimes^\nu [2]\ )$ $\quad (* = +, -, 0, \quad \nu \in \mathbb{N} \cup \{0\})$ *such that*

$$\Delta_{p+\nu}(\ K_+ \otimes^\nu [2]\ ) - \Delta_{p+\nu}(\ K_- \otimes^\nu [2]\ ) = (t + (-1)^{\nu+1}) \cdot \Delta_{p+\nu}(\ K_0 \otimes^\nu [2]\ ),$$

where [2] *denotes the empty knot* [2].

11. A REMARK ON $\mathbb{Z}[t, t^{-1}]$-CASE

Some of our results on polynomial invariants could be extended to the case where the word, '$\mathbb{Q}[t, t^{-1}]$-balanced class,' is replaced with the word, '($\mathbb{Z}[t, t^{-1}]$-balanced of) an element of $\mathbb{Z}[t, t^{-1}]$'. However, we must take care of the following Theorem 11.1.

Two polynomials $f(t), g(t) \in \mathbb{Z}[t, t^{-1}]$ are said to be $\mathbb{Z}[t, t^{-1}]$*-balanced* if there is an integer n such that $f(t) = \pm t^n \cdot g(t)$.

Theorem 11.1. *There is a smooth oriented codimension two closed submanifold* $K \subset S^{n+2}$ *with the following property.*
(1) *Any Seifert hypersurface of* K *satisfies the condition that the* p*-th betti number is equal to the* $(n + 1 - p)$*-th betti number* $(1 \leqq p \leqq n)$*.*
(2) *There are Seifert hypersurfaces* V *and* W *for* K *such that the* $\mathbb{Z}[t, t^{-1}]$*-balanced class of the determinant of a* p*-Alexander matrix of* V *and that of* W *are different for a non-negative integer* p *even if the linear map defined by a* $(p-1)$*-Alexander matrix associated with* V *(resp.* W*) is injective.*

Note. The $\mathbb{Z}[t, t^{-1}]$-balanced class of the determinant of a p-Alexander matrix of V (resp. W) is same as that of any p-Alexander matrix of V (resp. W) .

If the linear map defined by a $(p-1)$-Alexander matrix associated with V (resp. W) is injective, then that defined by any $(p-1)$-Alexander matrix associated with V (resp. W) is injective.

Because they do not depend on the choice of the basis of the homology groups of V (resp. W).

Proof. Let K be a closed oriented smooth 3-dimensional submanifold $\subset S^5$ such that the diffeomorphism type of K is represented by the following framed link: Take the (a,a)-torus link $\subset S^3$ $(a \in \mathbb{N} - \{1\})$. The framing of each component is zero. Then K is diffeomorphic to a homology sphere and, therefore, any Seifert hypersurface for K satisfies the condition (1) in Theorem 11.1. The following figure is the $a = 2$ case of the framed link.



We make two kinds of Seifert hypersurfaces V, W for K as follows.

The first. Regard $\mathbb{R}^5 = \mathbb{R}^4 \times \{t \in \mathbb{R}\}$. Regard the above framed link which represents K as a 4-manifold, too. Of course, this 4-manifold has a handle decomposition (a 4-dimensional 0-handle)$\cup$(a 4-dimensional 2-handle) $\cup$(a 4-dimensional 2-handle), which is defined by the framed link representation.

Suppose that the diffeomorphism type of a Seifert hypersurface V is this 4-manifold.

Suppose that V in $\mathbb{R}^5$ satisfies the following.

(1) The 4-dimensional 0-handle is embedded in $\mathbb{R}^4 \times \{t = 0\}$.

(2) One of the 4-dimensional 2-handles is embedded in $\mathbb{R}^4 \times \{t = 0\}$, call it h^2.

(3) The other of the 4-dimensional 2-handles is embedded in $\mathbb{R}^4 \times \{t \leqq 0\}$. Only the attached part is embedded in $\mathbb{R}^4 \times \{t = 0\}$. We can do this because the framing is zero.

Thus we obtain a Seifert hypersurface V for K.

Then a positive 2-Seifert matrix associated with V is $\begin{pmatrix} 0 & a \\ 0 & 0 \end{pmatrix}$. The negative 2-Seifert matrix associated with the positive 2-Seifert matrix is $\begin{pmatrix} 0 & 0 \\ -a & 0 \end{pmatrix}$. Hence the 2-Alexander matrix associated with these two matrices is $\begin{pmatrix} 0 & at \\ a & 0 \end{pmatrix}$. Its determinant is $-a^2 t$.

Note that the linear map defined by a 1-Alexander matrix associated with V is injective.

The second. Take the above Seifert hypersurface V. Suppose that a 5-dimensional 3-handle k^3 is embedded in $\mathbb{R}^4 \times \{t \geqq 0\}$. Attach k^3 along the 2-sphere embedded in V which includes the core of the above h^2. Suppose that only the attach part is embedded in $\mathbb{R}^4 \times \{t = 0\}$. By this surgery by k^3, V is changed into another Seifert hypersurface W for K. Then the framed link representation of W is as follows: Take the (a, a)-torus link $\subset S^3$. The framing of one component is zero. The other component is the dot circle (see [13] for the dot circle). Then W is a rational homology ball.

The following figure is the $a = 2$ case of W.



Hence the following hold. The positive 2-Seifert matrix associated with W is 'empty'. The negative 2-Seifert matrix associated with W is 'empty'. Hence the 2-Alexander matrix associated with W is 'empty'. Note that the $\mathbb{Z}[t, t^{-1}]$-balanced class of the determinant of the 2-Alexander matrix 'empty' is that of 1. Note that the linear map defined by a 1-Alexander matrix associated with W is injective. Note that the $\mathbb{Z}[t, t^{-1}]$-balanced class of the determinant of the 2-Alexander matrix 'empty' is NOT that of a^2t. (Recall that we define $a \in \mathbb{N} - \{1\}$.)

This completes the proof. □

12. Proof of Theorems in §8

We use the following proposition.

Proposition 12.1. *Let K be an n-dimensional closed oriented submanifold $\subset S^{n+2}$.*
(1) $S_{p+1}(K \otimes [2]) = (-1)^{(n-p)} S_p(K) \otimes S_0([2])$
(2) $N_{p+1}(K \otimes [2]) = (-1)^{(n-p)} N_p(K) \otimes N_0([2])$
(3) $S_0([2]) = (1)$.
(4) $N_0([2]) = (-1)$.
(5) $S_{\nu-1}(\otimes^\nu [2]) = (-1)^{\frac{(\nu-1)\nu}{2}}$.
Note that $\otimes^\nu[2]$ is a $(2\nu - 3)$-submanifold $\subset S^{2\nu-1}$.
(6) $S_{p+\nu}(K \otimes^\nu [2]) = (-1)^{(n-p)\nu + \frac{\nu(\nu-1)}{2}} \quad S_p(K)$
(7) $S_{p+2\mu}(K \otimes^\mu$ *(the Hopf link)*$) = (-1)^\mu S_p(K)$. *(Let $\nu = 2\mu$ in* (6).*)*

Let α, β be disjoint cycles of dimension p and q in S^{p+q+1}.

(8) $\mathrm{lk}(\alpha,\beta)=(-1)^{pq+1}\mathrm{lk}(\beta,\alpha)$.

By §6 of [9] we have (1)-(7). By page 541 of [14] we have (8).

Proof of Theorem 8.1. Take 1-links $K_+, K_- \subset S^3 = \partial B^4 \subset B^4$. The 1-links, K_+, K_-, differ by only one crossing-change in a 3-ball A trivially embedded in S^3 as shown below.



$A \cap K_+$



$A \cap K_-$

Take a Seifert hypersurface V_+ (resp. V_-) for the 1-link K_+ (resp. K_-) such that the submanifolds, V_+, V_-, differ only in the 3-ball $A \subset S^3$ as shown below.



$A \cap V_+$



$A \cap V_-$

Push V_+ (resp. V_-) into B^4, fixing $\partial V_+ = K_+$ (resp. $\partial V_- = K_-$), call the submanifold $\subset B^4$, V'_+ (resp. V'_-).

Suppose that the submanifolds, V'_+, V'_-, differ only in the 4-ball $A \times [0,1] \subset B^4$, where $A \times \{0\} = A \subset S^3$.

Note that $A \times [0,1]$ and $B^4 - ((\mathrm{Int}\ A) \times [0,1))$ are diffeomorphic to the 4-ball. The submanifold $A \times [0,1]$ (resp. $B^4 - ((\mathrm{Int}\ A) \times [0,1))$) is called B^4_I (resp. B^4_O). The submanifold $B^4_I \cap B^4_O$ is called $B^3_\cap$.

Let $B^2 = \{(x,y) \in \mathbb{R}^2 | x^2+y^2 \leqq 1\}$. Then there are smooth maps $f_+ : B^4 \to B^2$ and $f_- : B^4 \to B^2$ with the following properties. (Because: Use the Thom-Pontrjagin construction.)

(1) The submanifold $f_+^{-1}((0,0)) \subset B^4$ is V'_+.
The submanifold $f_-^{-1}((0,0)) \subset B^4$ is V'_-.

(2) $f_+|_{B^4_O} = f_-|_{B^4_O}$

Make the knot products $K_+ \otimes$ (the empty knot [2]) and $K_- \otimes$ (the empty knot [2]).

See Definition 5.1 and [8, 9]. The knot product $K_* \otimes$ (the empty knot [2]) is defined as follows ($* = +, -$). Take a smooth map $g : B^2 \to B^2$ such that $(r\cos\theta, r\sin\theta) \mapsto (r\cos 2\theta, r\sin 2\theta)$, where we use the standard polar coordinate.

Then $g|_{\partial B^2 = S^1} : S^1 \to S^1$ is the empty knot [2].
Let $M_* = \{(x, y) \in B^4 \times B^2 | f_*(x) - g(y) = (0, 0) \in B^2\}$.
The 3-dimensional closed oriented submanifold $\partial M_* \subset \partial(B^4 \times B^2) = \check{S}^5$ is the knot product $K_* \otimes [2]$.

Since $f_+|_{B^4_O} = f_-|_{B^4_O}$, we have $M_+ \cap (B^4_O \times B^2) = M_- \cap (B^4_O \times B^2)$. (They are identical as subsets $\subset B^4_O \times B^2$.) Therefore we have
$\partial M_+ \cap (\check{S}^5 \cap (B^4_O \times B^2)) = \partial M_- \cap (\check{S}^5 \cap (B^4_O \times B^2))$. (They are identical as submanifolds $\subset \check{S}^5 \cap (B^4_O \times B^2)$.)
Note that $\check{S}^5$ is the union of $\check{S}^5 \cap (B^4_O \times B^2)$ and $\check{S}^5 \cap (B^4_I \times B^2)$. Note that $\check{S}^5 \cap (B^4_O \times B^2)$ and $\check{S}^5 \cap (B^4_I \times B^2)$ are diffeomorphic to the 5-ball. $\check{S}^5 \cap (B^4_O \times B^2)$ (resp. $\check{S}^5 \cap (B^4_I \times B^2)$) is called $\check{B}^5_O$ (resp. $\check{B}^5_I$).

Note that the intersection of $\check{B}^5_O = \check{S}^5 \cap (B^4_O \times B^2)$ and $\check{B}^5_I = \check{S}^5 \cap (B^4_I \times B^2)$ is
$\check{S}^5 \cap (B^3_\cap \times B^2)$, is diffeomorphic to the standard 4-sphere, is called $\check{S}^4_\cap$.
Furthermore, the fact $f_+|_{B^4_O} = f_-|_{B^4_O}$ claims the following.
(1) $(K_+ \otimes [2]) \cap \check{B}^5_O = (K_- \otimes [2]) \cap \check{B}^5_O$. (They are identical as submanifolds $\subset \check{B}^5_O$.)
(2) $(K_+ \otimes [2]) \cap \check{S}^4_\cap = (K_- \otimes [2]) \cap \check{S}^4_\cap$. (They are identical as submanifolds $\subset \check{S}^4_\cap$.)

By [8, 9], the diffeomorphism type of $(K_+ \otimes [2]) \cap \check{B}^5_I$ (resp. $(K_- \otimes [2]) \cap \check{B}^5_I$) is the 2-fold branched cyclic covering space of the 3-ball A along the two arcs $K_+ \cap A$ (resp. $K_- \cap A$). It is the solid torus.

By [8, 9], the diffeomorphism type of $(K_+ \otimes [2]) \cap \check{S}^4_\cap$ (resp. $(K_- \otimes [2]) \cap \check{S}^4_\cap$) is the 2-fold branched cyclic covering space of the 2-sphere ∂A along the four points $K_+ \cap \partial A$ (resp. $K_- \cap \partial A$). It is the torus.

By [8, 9], we have the following.
(1) There is a Seifert hypersurface W_* for $K_* \otimes [2] \subset \check{S}^5$ $(* = +, -)$ which is diffeomorphic to the 2-fold branched covering space of B^4 along V'_*.
(2) Take the identity map $id : \check{S}^5_O \to \check{S}^5_O$. Then $id|_{\check{B}^5_O}$ is the identity map: $\check{B}^5_O \to \check{B}^5_O$. Furthermore, $id(W_+ \cap \check{B}^5_O) = W_- \cap \check{B}^5_O$.

When we push off $V_* \subset S^3$ into B^4, we can suppose the following.
Let $\beta : V_* \times [0, 1] \to B^4 (0 \leqq t \leqq 1)$ be the isotopy. Then $(A \times [0, 1]) \cap \beta(V_*, t)$ is diffeomorphic to $A \cap V_*$ for any t. Furthermore, $(A \cap V_*) \times [0, 1] = (A \times [0, 1]) \cap \beta(V_*, [0, 1])$. (They are identical as submanifolds $\subset B^4$.)

By [8, 9], the diffeomorphism type of $V'_+ \cap B^4_I$ and that of $V'_- \cap B^4_I$ are same and are diffeomorphic to $A \cap V_*$. Recall $B^4_I = A \times [0, 1]$. The submanifold $V'_+ \cap A \times \{0\}$ and $V'_- \cap A \times \{0\}$ are equivalent to the two arcs in A in the first figure in this Proof.

The diffeomorphism type of $W_+ \cap \check{B}^5_I$ and that of $W_- \cap \check{B}^5_I$ are same. By [8, 9], $W_* \cap \check{B}^5_I$ is diffeomorphic to the 2-fold branched covering space of B^4_I along $V'_* \cap B^4_I$ $(* = +, -)$.

Hence $W_* \cap \check{B}^5_I$ is diffeomorphic to the 4-ball for each case $* = +, * = -$. The intersection of the 4-ball $W_* \cap \check{B}^5_I$ and $\partial \check{B}^5_I = \check{S}^4_\cap$ is the solid torus whose boundary is $(K \otimes [2]) \cap \check{S}^4_\cap$.

Therefore we can regard $W_* \cap \check{B}^5_I$ as an embedded 4-dimensional 2-handle ($\subset \check{B}^5_I \subset \check{S}^5$) which is attached to $W_* \cap \check{B}^5_O$ along the solid torus $W_* \cap \partial \check{B}^5_O = W_* \cap \check{S}^4_\cap$. The diffeomorphism map of $* = +$ is same as that of $* = -$.

Since K_+ and K_- differ by only one crossing-change, there is a Seifert matrix $X_* = (x^*_{i,j})$ for K_* with the following property ($* = +.-$).

$$\left\{\begin{array}{l} x^+_{11} = 1, \quad x^-_{11} = 0 \\ x^+_{ij} = x^-_{ij} \quad \text{if } (i,j) \neq (1,1). \quad i,j \leqq \nu. \quad i,j \in \mathbb{N} - \{0\}. \end{array}\right.$$

By [8, 9] and Proposition 12.1, we can suppose that the $\nu \times \nu$-matrix $-X_* = (-x^*_{i,j})$ is a 2-Seifert matrix for $K_* \otimes [2]$.

Recall the definition of twist-moves in §4. By the above two paragraphs, $K_+ \otimes [2]$ and $K_- \otimes [2]$ differ by one twist-move.

This completes the proof. $\square$

Note. The above proof is the $m = 0, \nu = 1$ case of the Proof of Theorem 8.3.

Proof of Theorem 8.2. This proof is the $m = 0$ case of Proof of Theorem 8.3. $\square$

Proof of Theorem 8.3. We first prove the $\nu = 1$ case.

Take $(2m+1)$-dimensional oriented closed submanifolds $K_+, K_- \subset S^{2m+3} = \partial B^{2m+4} \subset B^{2m+4}$. The $(2m+1)$-dimensional oriented closed submanifolds, K_+, K_-, differ by only one twist-move in a $(2m+3)$-ball A trivially embedded in S^{2m+3}. See the left two figures in the upper half of Figure 4.5. Take a Seifert hypersurface V_+ (resp. V_-) for the $(2m+1)$-dimensional oriented closed submanifold K_+ (resp. K_-) such that the submanifolds, V_+, V_-, differ by only one twist-move in the $(2m+3)$-ball $A \subset S^{2m+3}$ and that $V_* \cap A$ is the $(2m+2)$-dimensional $(m+1)$-handle $h_*(* = +, -)$. See the definition of twist-moves in §4. Push V_+ (resp. V_-) into B^{2m+4}, fixing $\partial V_+ = K_+$ (resp. $\partial V_- = K_-$), call the submanifold $\subset B^{2m+4}$, V'_+ (resp. V'_-). Suppose that the submanifolds, V'_+, V'_-, differ only in the $(2m+4)$-ball $A \times [0,1] \subset B^{2m+4}$, where $A \times \{0\} = A \subset S^{2m+3}$. Note that $A \times [0,1]$ and $B^{2m+4} - ((\text{Int } A) \times [0,1))$ are diffeomorphic to the $(2m+4)$-ball. The submanifold $A \times [0,1]$ (resp. $B^{2m+4} - ((\text{Int } A) \times [0,1))$) is called B^{2m+4}_I (resp. B^{2m+4}_O). The submanifold $B^{2m+4}_I \cap B^{2m+4}_O$ is called $B^{2m+3}_\cap$. Let $B^2 = \{(x,y) \in \mathbb{R}^2 | x^2 + y^2 \leqq 1\}$. Then there are smooth maps $f_+ : B^{2m+4} \to B^2$ and $f_- : B^{2m+4} \to B^2$ with the following properties. (Because: Use the Thom-Pontrjagin construction.)

(1) The submanifold $f_+^{-1}((0,0)) \subset B^{2m+4}$ is V'_+.
    The submanifold $f_-^{-1}((0,0)) \subset B^{2m+4}$ is V'_-.

(2) $f_+|_{B^{2m+4}_O} = f_-|_{B^{2m+4}_O}$

Make the knot products $K_+ \otimes$ (the empty knot [2]) and $K_- \otimes$ (the empty knot [2]).

See Definition 5.1 and [8, 9]. The knot product $K_* \otimes$ (the empty knot [2]) is defined as follows ($* = +, -$). Take a smooth map $g : B^2 \to B^2$ such that
$(r\cos\theta, r\sin\theta) \mapsto (r\cos 2\theta, r\sin 2\theta)$, where we use the standard polar coordinate.
Then $g|_{\partial B^2 = S^1} : S^1 \to S^1$ is the empty knot [2].
Let $M_* = \{(x, y) \in B^{2m+4} \times B^2 | f_*(x) - g(y) = (0, 0) \in B^2\}$.
The $(2m+3)$-dimensional closed oriented submanifold $\partial M_* \subset \partial(B^{2m+4} \times B^2) = \check{S}^{2m+5}$ is the knot product $K_* \otimes [2]$.

Since $f_+|_{B_O^{2m+4}} = f_-|_{B_O^{2m+4}}$, we have $M_+ \cap (B_O^{2m+4} \times B^2) = M_- \cap (B_O^{2m+4} \times B^2)$. (They are identical as subsets $\subset B_O^{2m+4} \times B^2$.) Therefore we have
$\partial M_+ \cap (\check{S}^{2m+5} \cap (B_O^{2m+4} \times B^2)) = \partial M_- \cap (\check{S}^{2m+5} \cap (B_O^{2m+4} \times B^2))$. (They are identical as submanifolds $\subset \check{S}^{2m+5} \cap (B_O^{2m+4} \times B^2)$.)
Note that $\check{S}^{2m+5}$ is the union of $\check{S}^{2m+5} \cap (B_O^{2m+4} \times B^2)$ and $\check{S}^{2m+5} \cap (B_I^{2m+4} \times B^2)$. Note that $\check{S}^{2m+5} \cap (B_O^{2m+4} \times B^2)$ and $\check{S}^{2m+5} \cap (B_I^{2m+4} \times B^2)$ are diffeomorphic to the $(2m+5)$-ball. $\check{S}^{2m+5} \cap (B_O^{2m+4} \times B^2)$ (resp. $\check{S}^{2m+5} \cap (B_I^{2m+4} \times B^2)$) is called $\check{B}_O^{2m+5}$ (resp. $\check{B}_I^{2m+5}$).

Note that the intersection of $\check{B}_O^{2m+5} = \check{S}^{2m+5} \cap (B_O^{2m+4} \times B^2)$ and
$\check{B}_I^{2m+5} = \check{S}^{2m+5} \cap (B_I^{2m+4} \times B^2)$ is $\check{S}^{2m+5} \cap (B_\cap^{2m+3} \times B^2)$, is diffeomorphic to the standard $(2m+4)$-sphere, is called $\check{S}_\cap^{2m+4}$.
Furthermore, the fact $f_+|_{B_O^{2m+4}} = f_-|_{B_O^{2m+4}}$ claims the following.
(1) $(K_+ \otimes [2]) \cap \check{B}_O^{2m+5} = (K_- \otimes [2]) \cap \check{B}_O^{2m+5}$. (They are identical as submanifolds $\subset \check{B}_O^{2m+5}$.)
(2) $(K_+ \otimes [2]) \cap \check{S}_\cap^{2m+4} = (K_- \otimes [2]) \cap \check{S}_\cap^{2m+4}$. (They are identical as submanifolds $\subset \check{S}_\cap^{2m+4}$.)

By [8, 9], the diffeomorphism type of $(K_+ \otimes [2]) \cap \check{B}_I^{2m+5}$ (resp. $(K_- \otimes [2]) \cap \check{B}_I^{2m+5}$) is the 2-fold branched cyclic covering space of the $(2m+3)$-ball A along $K_+ \cap A$ (resp. $K_- \cap A$). It is diffeomorphic to $S^{m+1} \times B^{m+2}$.

By [8, 9], the diffeomorphism type of $(K_+ \otimes [2]) \cap \check{S}_\cap^{2m+4}$ (resp. $(K_- \otimes [2]) \cap \check{S}_\cap^{2m+4}$) is the 2-fold branched cyclic covering space of the $(2m+2)$-sphere ∂A along $K_+ \cap \partial A$ (resp. $K_- \cap \partial A$). It is diffeomorphic to $S^{m+1} \times S^{m+1}$.

By [8, 9], we have the following.
(1) There is a Seifert hypersurface W_* for $K_* \otimes [2] \subset \check{S}^{2m+5}$ ($* = +, -$) which is diffeomorphic to the 2-fold branched covering space of B^{2m+4} along V'_*.
(2) Take the identity map $id : \check{S}_O^{2m+5} \to \check{S}_O^{2m+5}$. Then $id|_{\check{B}_O^{2m+5}}$ is the identity map $\check{B}_O^{2m+5} \to \check{B}_O^{2m+5}$. Furthermore, $id(W_+ \cap \check{B}_O^{2m+5}) = W_- \cap \check{B}_O^{2m+5}$.

When we push off $V_* \subset S^{2m+3}$ into B^{2m+4}, we can suppose the following.
Let $\beta : V_* \times [0, 1] \to B^{2m+4} (0 \leqq t \leqq 1)$ be the isotopy. Then $(A \times [0, 1]) \cap \beta(V_*, t)$ is diffeomorphic to $A \cap V_*$ for any t. Furthermore, $(A \cap V_*) \times [0, 1] = (A \times [0, 1]) \cap \beta(V_*, [0, 1])$. (They are identical as submanifolds $\subset B^{2m+4}$.)

By [8, 9], the diffeomorphism type of $V'_+ \cap B_I^{2m+4}$ and that of $V'_- \cap B_I^{2m+4}$ are same and are diffeomorphic to $A \cap V_*$. Recall $B_I^{2m+4} = A \times [0,1]$. Recall that the submanifold $V'_+ \cap A \times \{0\}$ (resp. $V'_- \cap A \times \{0\}$) in $A \times \{0\}$ is $K_+ \cap A$ (resp. $K_- \cap A$) in A.

The diffeomorphism type of $W_+ \cap \check{B}_I^{2m+5}$ and that of $W_- \cap \check{B}_I^{2m+5}$ are same. By [8, 9], $W_* \cap \check{B}_I^{2m+5}$ is diffeomorphic to the 2-fold branched covering space of B_I^{2m+4} along $V'_* \cap B_I^{2m+4}$ ($* = +, -$). Hence $W_* \cap \check{B}_I^{2m+5}$ is diffeomorphic to the $(2m+4)$-ball for each case $* = +, * = -$. The intersection of the $(2m+4)$-ball and $\partial \check{B}_I^{2m+5} = \check{S}_\cap^{2m+4}$ is diffeomorphic to $S^{m+1} \times B^{m+2}$ whose boundary is $(K_* \otimes [2]) \cap \check{S}_\cap^{2m+4}$.

Therefore we can regard $W_* \cap \check{B}_I^{2m+5}$ as an embedded $(2m+4)$-dimensional $(m+2)$-handle ($\subset \check{B}_I^{2m+5} \subset \check{S}^{2m+5}$) which is attached to $W_* \cap \check{B}_O^{2m+5}$ along $W_* \cap \partial \check{B}_O^{2m+5}$ $= W_* \cap \check{S}_\cap^{2m+4}$. The diffeomorphism map of $* = +$ is same as that of $* = -$.

Since K_+ and K_- differ by only one crossing-change, there is an $(m+1)$-Seifert matrix $X_* = (x^*_{i,j})$ for K_* with the following property ($* = +.-$).

$$\begin{cases} x^+_{11} = 1, \quad x^-_{11} = 0 \\ x^+_{ij} = x^-_{ij} \quad \text{if } (i,j) \neq (1,1). \quad i, j \leqq \nu. \quad i, j \in \mathbb{N} - \{0\}. \end{cases}$$

By [8, 9] and Proposition 12.1, we can suppose that the $\nu \times \nu$-matrix $-X_* = (-x^*_{i,j})$ is an $(m+2)$-Seifert matrix for $K_* \otimes [2]$.

Recall the definition of twist-moves in §4. By the above two paragraphs, $K_+ \otimes [2]$ and $K_- \otimes [2]$ differs by one twist-move.

Next we prove the $\nu > 1$ case.

If the '$2m = 2\alpha$ and $\nu = 1$' case is true, then the '$2m = 2\alpha + 2$ and $\nu = 1$' case is true. Hence '$2m = 2\alpha$ and $\nu = 2$' case is true. By the induction, Theorem 8.3 is true. □

Here, we prove the following Proposition 12.2, which is used in §14.

Suppose that two $(2m+1)$-dimensional oriented closed submanifolds $\subset S^{2m+3}$, K_+ and K_-, differ by one twist-move. By Theorem 8.3, the $(2m+2\nu+1)$-submanifolds $\subset S^{2m+2\nu+3}$, $K_+ \otimes^\nu [2]$ and $K_- \otimes^\nu [2]$, differ by one twist-move in a $(2m+2\nu+3)$-ball A. Then there is a unique closed oriented $(2m+2\nu+1)$-submanifold $K_0^\otimes \subset S^{2m+2\nu+3}$ such that a triple $(K_+ \otimes^\nu [2],\ K_- \otimes^\nu [2],\ K_0^\otimes)$ is related twist-move in A. Note that the equivalence class of the submanifold $K_0^\otimes \subset S^{2m+2\nu+3}$ is determined uniquely. Note that we have the following. Take $W_+ = W_-$ as in the $(2m+2\nu+1)$ case of Proof of Theorem 8.3. Then $\partial(W_+ - \text{Int}B^{2m+2\nu+3}) = \partial(W_- - \text{Int}B^{2m+2\nu+3})$ in $S^{2m+2\nu+3}$ is $K_0^\otimes \subset S^{2m+2\nu+3}$. Then we have the following.

Proposition 12.2. *The* $(2m+2\nu+1)$*-submanifolds* $\subset S^{2m+2\nu+3}$*,* $K_0^\otimes$ *and* $K_0 \otimes^\nu$ (*the empty knot* $[2]$)*, are equivalent.*

Proof of Theorem 12.2. It suffices to prove the $\nu = 1$ case.

Take $M_+ \cap (S^{2m+5} \cap (B^{2m+4}_O \times B^2)) = M_- \cap (S^{2m+5} \cap (B^{2m+4}_O \times B^2))$ in the above Proof of Theorem 8.3. By the construction, the submanifold $M_* \cap \partial(S^{2m+5} \cap (B^{2m+4}_O \times B^2))$ in $\partial(S^{2m+5} \cap (B^{2m+4}_O \times B^2))$ $(* = +, -)$ is $K_0^{\otimes}$ and $K_0 \otimes^\nu [2] \subset S^{2m+2\nu+3}$. $\square$

Proof of Theorem 8.4.

Proof of Theorem 8.4.(1). Take the 1-knot K in the following figure.



Twist in the shaded part so that its Seifert matrix is $\begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}$.

By [32], K is a nontrivial knot. Note that the unknotting number of K is one.

By [8, 9] and Proposition 12.1, the $(2\nu+1)$-submanifold $K \otimes^\nu [2] \subset S^{2\nu+3}$ has a Seifert hypersurface V with the following conditions.

(1) V has a handle decomposition
(a $(2\nu+2)$-dimensional 0-handle)$\cup$(($2\nu+2)$-dimensional $(\nu+1)$-handles), where there may be no $(2\nu+2)$-dimensional $(\nu+1)$-handle.

(2) A Seifert matrix S associated with V is $\begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}$ or $\begin{pmatrix} -1 & -1 \\ 0 & 0 \end{pmatrix}$.

Note that $S + (-1)^{\nu+1}({}^tS)$ represents the intersection product of $H_{\nu+1}(V;\mathbb{Z})/(\mathrm{Tor})$. Since the determinant of $S + (-1)^{\nu+1}({}^tS)$ is $+1$ or -1, $\partial V = K \otimes^\nu [2]$ is a homology sphere. Recall $\nu \geqq 2$. By (1) right above, $\pi_1 \partial V = 1$. By [28], ∂V is PL homeomorphic to the standard sphere. Hence the $(2\nu+1)$-submanifold $K \otimes^\nu [2]$ is a spherical knot. By (1) right above, the $(2\nu+1)$-knot $K \otimes^\nu [2]$ is a simple knot.

Recall that the trivial $(2\nu+1)$-knot has Seifert matrices, $\begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}$ and $\begin{pmatrix} -1 & -1 \\ 0 & 0 \end{pmatrix}$.

By [16], the $(2\nu+1)$-knot $K \otimes^\nu [2]$ is equivalent to the trivial knot. $\square$

Proof of Theorem 8.4.(2). Take the above nontrivial 1-knot K. Take the knot-sum $K\sharp K$. By [6, 25] and the van Kampen's theorem, $K\sharp K$ is nontrivial. By [27], the unknotting number of $K\sharp K$ is two.

Note $(K\sharp K)\otimes^\nu [2]$ is the trivial knot.

$K\sharp K$ is an example which we want. $\square$

Proof of Theorem 8.4.(3). The pair of K and $K\sharp K$ is an example which we want. Because: K is a prime knot and $K\sharp K$ is not. Hence K is not equivalent to $K\sharp K$. $(K\sharp K)\otimes^\nu [2]$ and $K\otimes^\nu [2]$ are trivial knots and hence equivalent spherical knots. $\square$

Proof of Theorem 8.4.(4). Let A be the trefoil knot (and hence a nontrivial 1-knot). By [6, 25] and the van Kampen's theorem, $A\sharp K$ is nontrivial. By [27], the unknotting number of $A\sharp K$ is two. By [8, 9] and [16], $(A\sharp K)\otimes^\mu$(the Hopf link) and $A\otimes^\mu$(the Hopf link) are equivalent and nontrivial . By Theorem 8.2, $A\otimes^\mu$(the Hopf link) is obtained from the trivial knot by a twist-move. Hence $(A\sharp K)\otimes^\mu$(the Hopf link) is obtained from the trivial knot by a twist-move. Hence $A\sharp K$ is an example which we want. $\square$

Proof of Theorem 8.4.(5). The pair of A and $A\sharp K$ is an example which we want. $\square$

Proof of $(*)$ in Note 8.5. Take the nontrivial 1-knot K in Proof of Theorem 8.4. The diffeomorphism type of $K\otimes[2]$ is the 3-manifold which is the double branched covering space of S^3 along K (see [8, 9].) It is not diffeomorphic to the standard 3-sphere (see [18, 29].) K is an example which we want. $\square$

In order to prove Theorem 8.6, we prove some propositions and a theorem.

Proposition 12.3. *Let $l\in\mathbb{N}$. Let K be a simple $(2l+1)$-knot. Then $K\otimes$ (the Hopf link) is a simple knot.*

Proof of Proposition 12.3. By [8, 9] and Proposition 12.1, $K\otimes$(the Hopf link) satisfies the following.

(1) A Seifert hypersurface has a handle decomposition of one $(2l+6)$-dimensional 0-handle and $(2l+6)$-dimensional $(l+3)$-handles, where there may not be $(2l+6)$-dimensional $(l+3)$-handles.

(2) A $(l+3)$-Seifert matrix Y associated with the above Seifert hypersurface is $(-1)\times$ (a $(l+1)$-Seifert matrix of K).

By these (1) and (2) and [28], $K\otimes$ (the Hopf link) is a spherical knot and a simple knot. $\square$

Proposition 12.4. *Let $p\in\mathbb{N}$. Suppose that K is a simple $(2p+5)$-knot and that, if $2p+5=7$, the signature of K is a multiple of 16. Then there is a simple $(2p+1)$-knot A with the following properties.*

(i) *K is equivalent to $A\otimes$ (the Hopf link).*

(ii) *If* X *is a* $(p+3)$*-Seifert matrix of* K*, then* $-X$ *is a* $(p+1)$*-Seifert matrix of* A*.*
(iii) *The equivalence class of such a knot is unique.*

Proof of Proposition 12.4. Take a $(p+3)$-Seifert matrix X of K. By [16] and Propsotion 12.1, there is a simple $(2p+1)$-knot A such that a $(p+1)$-Seifert matrix is the matrix $-X$. By Propsotion 12.1 and 12.3, $A \otimes$ (the Hopf link) is a simple $(2p+5)$-knot such that a $(p+2)$-Seifert matrix is the matrix X. By [16], $A \otimes$ (the Hopf link) is equivalent to K. By [16], (iii) holds. $\square$

Proposition 12.5. *Let* K *be a 1-knot. Let* $\mu \in \mathbb{N}$*. Then* $K \otimes^{\mu}$ (*the Hopf link*) *is a* $(4\mu+1)$*-dimensional simple knot* (*and hence a spherical knot*)*.*

Proof of Proposition 12.5. By [8, 9] and Proposition 12.1, $K \otimes^{\mu}$ (the Hopf link) satisfies the following.
(1) A Seifert hypersurface has a handle decomposition of one $(4\mu+2)$-dimensional 0-handle and $(4\mu+2)$-dimensional $(2\mu+1)$-handles, where there may not be $(2\mu+1)$-handles.
(2) Let Y be a Seifert matrix of K. A Seifert matrix associated with the above Seifert hypersurface is Y or $-Y$.

By (2) and the fact that K is diffeomorphic to the single circle, det $(Y - {}^tY)$ is ± 1. By this fact and the above (1) (2) and [28], $K \otimes^{\mu}$ (the Hopf link) is a spherical knot and a simple knot. $\square$

Proposition 12.6. *Let* K *be a simple 5-knot. Then there is a 1-knot* A *such that* K *is equivalent to* $A \otimes$ (*the Hopf link*)*.*

Proof of Proposition 12.6. Take a 3-Seifert matrix X of K. By using a Seifert surface, there is a 1-knot A such that a Seifert matrix is $-X$. By Propsotion 12.1 and 12.5, $A \otimes$ (the Hopf link) is a simple 5-knot such that a 3-Seifert matrix is X. By [16], $A \otimes$ (the Hopf link) is equivalent to K. $\square$

Note. The equivalence class of A is not unique. There are countably infinitely many equivalence types of 1-knots of this property. Because: Use the nontrivial knot K in Proof of Theorem 8.4. Take knot-sums.

Proposition 12.7. *Let* $p \in \mathbb{N}$*. Let* K *be a simple* $(2p+1)$*-knot. Let* J *be a* $(2p+1)$*-submanifold in* S^{2p+3} *such that* J *is obtained from* K *by one twist-move. Then* J *is a simple knot.*

Proof of Proposition 12.7. By the definition of twist-moves, there is a $(2p+3)$-ball B trivially embedded in S^{2p+3} in which this twist-move is carried out.

By using the Meyer-Vietoris exact sequence and the van Kampen's theorem, J is PL homeomorphic to the standard sphere.

Let $N(J)$ (resp. $N(K)$) be the tubular neighborhood of J (resp. $N(K)$) in S^{2p+3}. Then we have
$S^{2p+3}-\mathrm{Int}N(K) = (S^{2p+3}-\mathrm{Int}N(K)-\mathrm{Int}B) \cup$ (a $(2p+3)$-dimensional $(p+2)$-handle).
$S^{2p+3}-\mathrm{Int}N(J) = (S^{2p+3}-\mathrm{Int}N(J)-\mathrm{Int}B) \cup$ (a $(2p+3)$-dimensional $(p+2)$-handle).

By the definition of twist-moves, $S^{2p+3}-\mathrm{Int}N(K)-\mathrm{Int}B = S^{2p+3}-\mathrm{Int}N(J)-\mathrm{Int}B$.

Since K is a simple knot, $\pi_1(S^{2p+3}-\mathrm{Int}N(K)) = \mathbb{Z}$. Use the van Kampen's theorem for the above unions. Hence $\pi_1(S^{2p+3}-\mathrm{Int}N(J)) = \mathbb{Z}$.

By using the Thom-Pontrjagin construction, we can prove that there is a Seifert hypersurface V_K for K and V_J for J with the following properties.
(1) $V_K \cap B$ (resp. $V_J \cap B$) is a $(2p+2)$-dimensional $(p+1)$-handle that is attached to ∂B as explained in the definition of the twist-moves in §4.
(2) $V_K \cap (S^{2p+3}-\mathrm{Int}\ B) = V_J \cap (S^{2p+3}-\ \mathrm{Int}\ B)$.

Hence we have the following. Let $i \in \mathbb{N}$ and $i \leqq p$. there is an i-Seifert matrix X_{iK} (resp. X_{iJ}) for K (resp. J) such that $X_{iK} = X_{iJ}$. Consider the homology groups, the homotopy groups, and the fundamental group of the infinite cyclic covering space for K (resp. J). Hence J is a simple knot. □

We prove the following theorem.

Theorem 12.8. *Let $n \in \mathbb{N}\cup\{0\}$. Let K be an n-dimensional closed oriented submanifold $\subset S^{n+2}$. Take a map*

$$K \mapsto K \otimes (\textit{the Hopf link})$$

from the set of n-dimensional closed oriented submanifolds $\subset S^{n+2}$ to the set of n-dimensional closed oriented submanifolds $\subset S^{n+6}$.

Then we have the following.
(1) *Let K be a simple $(2l+1)$-knot ($l \geqq 2$, $l \in \mathbb{N}$). That is, suppose that the domain of the map is the set of simple $(2l+1)$-knots. Then the image of the map is the set of simple $(2l+5)$-knots. Furthermore, the map*

$$\{\textit{simple } (2l+1)\textit{-knots}\} \to \{\textit{simple } (2l+5)\textit{-knots}\}.$$
$$K \mapsto K \otimes (\textit{the Hopf link})$$

is a one-to-one map.
(2) *Let K be a simple 3-knot. That is, suppose that the domain of the map is the set of simple 3-knots. Then the image of the map is included in the set of simple 7-knots. Furthermore, the map*

$$\{\textit{simple 3-knots}\} \to \{\textit{simple 7-knots}\}.$$
$$K \mapsto K \otimes (\textit{the Hopf link})$$

is injective but not onto.
(3) *Let K be a 1-knot. That is, suppose that the domain of the map is the set of 1-knots. Then the image of the map is the set of simple 5-knots. Furthermore, the map*

$$\{1\text{-}knots\} \to \{simple\ 5\text{-}knots\}.$$
$$K \mapsto K \otimes (the\ Hopf\ link)$$

is onto but not injective. The inverse image of any element by this map is an infinite set.

Problem 12.9. What happens if we define the domain is another set in Theorem 12.8?

Proof of Theorem 12.8. Proposition 12.3, 12.4 imply Theorem 12.8.(1).

There is a simple 7-knot with the following property $(*)$:
$(*)$ The signature is a multiple of 8 but not a multiple of 16.
There is not a simple 3-knot with the above property $(*)$. See [16].

By these facts and Proposition 12.3, 12.4, Theorem 12.8.(2) holds.

Theorem 12.8.(3) follows from Proposition 12.5, 12.6, Note to Proposition 12.6, and Theorem 8.4. ☐

Proof of Theorem 8.6. By Theorem 12.8, K is a simple knot. By Proposition 12.7, J is a simple knot.

Then there is a $(p+3)$-Seifert matrix X (resp. Y) for J (resp. K) with the following properties.
(1) X and Y are $c \times c$ matrices for a natural number c.
(2) Let x_{ij} denote (i,j)-element of X. Let y_{ij} denote (i,j)-element of Y. There is a natural number $a \leqq c$ such that $\begin{cases} x_{ij} = y_{ij} - 1 & \text{if } (i,j) = (a,a) \\ x_{ij} = y_{ij} & \text{if } (i,j) \neq (a,a). \end{cases}$
(3) X and Y are not S-equivalent. See [16] for S-equivalent.

Note that we can take Seifert matrices which satisfy the above conditions. If necessarily, carry out surgery on a Seifert hypersurface by handles embedded in S^{2p+7}.

By Propsoition 12.1 and Theorem 12.8, there are simple $(2p+1)$-knots K' and J' with the following properties.
(i) J is equivalent to $J' \otimes$ (the Hopf link). $-X$ is a $(p+1)$-Seifert matrix for J'.
(ii) K is equivalent to $K' \otimes$ (the Hopf link). $-Y$ is a $(p+1)$-Seifert matrix for K'.
(iii) The equivalence class of such a knot is unique.

Therefore we can make a $(2p+1)$-dimensional simple knot $\bar{J}$ (resp. $\bar{K}$) with the following properties.
(I) A handle decomposition of a Seifert hypersurface is a set of a $(2p+2)$-dimensional 0-handle and $(2p+2)$-dimensional $(p+1)$-handles, where there may not be a $(2p+2)$-dimensional $(p+1)$-handle.
(II) A Seifert matrix associated with this Seifert hypersurface is $-X$ (resp. $-Y$).
(III) $\bar{J}$ and $\bar{K}$ differ by a single twist-move and are nonequivalent.

Because: Since J (resp. K) is PL homeomorphic to the standard sphere, we can realize (I) (II)(III) by using $(p+1, p+1)$-pass-moves.

By [16], simple $(2p+1)$-dimensional spherical knot $\bar{J}$ (resp. $\bar{K}$) is equivalent to J' (resp. K').

This completes the proof. $\square$

13. Proof of Theorems in §9

Proof of Theorem 9.1. There is a Seifert matrix X (resp. Y) for the 1-knot J (resp. K).

(1) X and Y are $c \times c$ matrices for a natural number c.

(2) Let x_{ij} denote (i,j)-element of X. Let y_{ij} denote (i,j)-element of Y. There are natural numbers $a, b \leqq c$ such that $a \neq b$ and that $\begin{cases} x_{ij} = y_{ij} - 1 & \text{if } (i,j) = (a,b) \\ x_{ij} = y_{ij} & \text{if } (i,j) \neq (a,b) \text{ and if } (i,j) \neq (b,a). \end{cases}$

The (b,a)-element is determined by the (a,b)-element.

Note that we can take Seifert matrices which satisfy the above conditions. If necessarily, carry out surgeries on Seifert hypersurfaces by 2-dimensional 1-handles embedded in S^3.

By Proposition 12.5, $J \otimes^{\mu}$ (the Hopf link) (resp. $K \otimes^{\mu}$ (the Hopf link)) is a spherical knot and is a simple knot.

We can make a $(4\mu+1)$-dimensional spherical knot J' (resp. K') with the following properties.

(1)A handle decomposition of a Seifert hyper surface is a set of a $(4\mu+2)$-dimensional $(2\mu+1)$-handle and $(4\mu+2)$-dimensional $(2\mu+1)$-handles, where there may not be $(4\mu+2)$-dimensional $(2\mu+1)$-handles.

(2) A Seifert matrix associated with the above Seifert hypersurface is $(-1)^{\mu}X$ (resp. $(-1)^{\mu}Y$).

(3)J' is obtained from K' by a $(2\mu+1, 2\mu+1)$-pass-move.

Because: Since J (resp. K) is diffeomorphic to the single circle, we can realize (1)(2)(3) by using $(2\mu+1, 2\mu+1)$-pass-moves.

By [16], simple $(4\mu+1)$-dimensional spherical knot J' (resp. K') is equivalent to $J \otimes^{\mu}$ (the Hopf link) (resp. $K \otimes^{\mu}$ (the Hopf link)).

By the construction of J' (resp. K'), $J \otimes^{\mu}$ (the Hopf link) and $K \otimes^{\mu}$ (the Hopf link) differ by one $(2\mu+1, 2\mu+1)$-pass-move. $\square$

Proof of Theorem 9.3. The pass-move does not change diffeomorphism type of submanifolds. However, $J \otimes^{\mu}$ [2] and $K \otimes^{\mu}$ [2] do not have the same homeomorphism type in general by [8, 9]. Example: (The trivial 1-knot)$\otimes$[2] and (the trefoil knot)$\otimes$[2]. $\square$

Proof of Theorem 9.5. Note $l \geqq 1$. There is a Seifert matrix X (resp. Y) for the simple $(2l+1)$-knot J (resp. K) with the following properties.
(1) X and Y are $c \times c$ matrices for a natural number c.
(2) Let x_{ij} denote (i,j)-element of X. Let y_{ij} denote (i,j)-element of Y. There are integers $a, b \leqq c$ such that $a \neq b$. We have $\begin{cases} x_{ij} = y_{ij} - 1 & \text{if } (i,j) = (a,b) \\ x_{ij} = y_{ij} & \text{if } (i,j) \neq (a,b) \text{ and if } (i,j) \neq (b,a) \end{cases}$
The (b,a)-element is determined by the (a,b)-element.

Note that we can take Seifert matrices which satisfy the above conditions. If necessarily, carry out surgeries on Seifert hypersurfaces by $(2l+2)$-dimensional $(l+1)$-handles embedded in S^{2l+3}.

By Proposition 12.3, $J \otimes^{\mu}$ (the Hopf link) (resp. $K \otimes^{\mu}$ (the Hopf link)) is a spherical knot and a simple knot.

We can make a $(2l+4\mu+1)$-dimensional spherical knot J' (resp. K') with the following properties.
(1) A handle decomposition of a Seifert hypersurface is a set of a $(2l+4\mu+2)$-dimensional 0-handle and $(2l+4\mu+2)$-dimensional $(l+2\mu+1)$-handles.
(2) A Seifert matrix associated with the above Seifert hypersurface is $(-1)^{\mu}X$ (resp. $(-1)^{\mu}Y$).
(3) J' is obtained from K' by a $(l+2\mu+1, l+2\mu+1)$-pass-move.
Because: Since J (resp. K) is PL homeomorphic to the standard sphere, we can realize (1)(2)(3) by using $(l+2\mu+1, l+2\mu+1)$-pass-moves.

By [16], simple $(2l+4\mu+1)$-dimensional spherical knot J' (resp. K') is equivalent to $J \otimes^{\mu}$ (the Hopf link) (resp. $K \otimes^{\mu}$ (the Hopf link)).

By the construction of J' (resp. K'), $J \otimes^{\mu}$ (the Hopf link) and $K \otimes^{\mu}$ (the Hopf link) differ by one $(l+2\mu+1, l+2\mu+1)$-pass-move. □

Proof of Theorem 9.8. Take the nontrivial 1-knot K in the figure in Proof of Theorem 8.4. Note that K is obtained from the trivial knot by one pass-move. □

Proof of $(*)$ in Note 9.10. It is same as Proof of $(*)$ in Note 8.5. □

Proof of Theorem 9.11. We need a proposition.

Proposition 13.1. *Let $p \in \mathbb{N}$. Let K be a simple $(2p+1)$-knot. Let J be a $(2p+1)$-submanifold in S^{2p+3} such that J is obtained from K by one $(p+1, p+1)$-pass-move. Then J is a simple knot.*

Proof of Proposition 13.1. By the definition of the $(p+1, p+1)$-pass-move, J is a spherical $(2p+1)$-knot.

See the following figure. We can take two copies of the $(p+1)$-sphere Y_1 and Y_2 in $B^{2p+3} \subset S^{2p+3}$ with the following property. Y_1 (resp. Y_2) is embedded trivially. The linking number of Y_1 and Y_2 is one. Carry out surgeries along two $(p+1)$-spheres Y_1

and Y_2 by two $(2p+3)$-dimensional $(p+2)$-handles with the trivial framing on B^{2p+3}. Then B^{2p+3} becomes the $(2p+3)$-ball again and S^{2p+3} becomes the $(2p+3)$-sphere again. Furthermore, the $(p+1, p+1)$-pass-move in the $(2p+3)$-ball B^{2p+3} is done.

Since $p \geqq 1$ holds and K is a simple knot, $\pi_i(S^{2p+3} - N(K)) = \pi_i(S^{2p+3} - N(K))$ for $1 \leqq i \leqq p$). (Use the van Kampen's theorem and the Meyer-Vietoris theorem on the complements and the infinite cyclic covering spaces.) Therefore J is a simple knot. This completes the proof of Proposition 13.1. $\square$



By Theorem 12.8, K is a simple knot. By Proposition 13.1, J is a simple knot. By Proposition 12.6 and Theorem 12.8, there is a 1-knot J' to satisfy (i).

By [19], $K'\otimes^\mu$(the Hopf link) and $J'\otimes^\mu$(the Hopf link) have the same Arf invariant.

By [8, 9], a $(2\mu+1)$-Seifert matrix of $\begin{cases} K'\otimes^\mu(\text{the Hopf link}) \\ J'\otimes^\mu(\text{the Hopf link}) \end{cases}$

is $(\pm 1)\times$ a Seifert matrix of $\begin{cases} K' \\ J' \end{cases}$. Hence Arf (K') =Arf(J').

By Theorem 3.2, K' is pass-move equivalent to J'. Hence J' satisfies (i) and (ii). This completes the proof of Theorem 9.11. $\square$

Note. Proof of Note to Theorem 9.11. Use the nontrivial knot K in Proof of Theorem 8.4. Take a knot-sum as many times as we need.

Proof of Theorem 9.12. By Theorem 12.8, K is a simple knot. By Proposition 12.7, J is a simple knot.

Then there is a $(p+3)$-Seifert matrix X (resp. Y) for J (resp. K) with the following properties.
(1) X and Y are $c \times c$ matrices for a natural number c.
(2) Let x_{ij} denote (i,j)-element of X. Let y_{ij} denote (i,j)-element of Y. There are integers $a, b \leqq c$ such that $a \neq b$. We have $\begin{cases} x_{ij} = y_{ij} - 1 & \text{if } (i,j) = (a,b) \\ x_{ij} = y_{ij} & \text{if } (i,j) \neq (a,b) \text{ and if } (i,j) \neq (b,a). \end{cases}$
The (b,a)-element is determined by the (a,b)-element.
(3) X and Y are not S-equivalent. See [16] for S-equivalent.

Note that we can take Seifert matrices which satisfy the above conditions. If necessarily, carry out surgeries on Seifert hypersurfaces by handles embedded in S^{2p+7}.

By Proposition 12.4 and Theorem 12.8, there are simple $(2p+1)$-knots K' and J' with the following properties.
(i) J is equivalent to $J' \otimes$ (the Hopf link). $-X$ is a $(p+1)$-Seifert matrix for J'.
(ii) K is equivalent to $K' \otimes$ (the Hopf link). $-Y$ is a $(p+1)$-Seifert matrix for K'.
(iii) The equivalence class of such a knot is unique.

Therefore we can make a $(2p+1)$-dimensional simple knot $\bar{J}$ (resp. $\bar{K}$) with the following properties.
(I) A handle decomposition of a Seifert hypersurface is a set of a $(2p+2)$-dimensional 0-handle and $(2p+2)$-dimensional $(p+1)$-handles, where there may not be a $(2p+2)$-dimensional $(p+1)$-handle.
(II) A Seifert matrix associated with the above Seifert hypersurface is $-X$ (resp. $-Y$).
(III) $\bar{J}$ and $\bar{K}$ differ by a single $(p+1, p+1)$-pass-move and are nonequivalent.
Because: Since J (resp. K) is PL homeomorphic to the standard sphere, we can realize (I) (II)(III) by using $(p+1, p+1)$-pass-moves.

By [16], simple $(2p+1)$-dimensional spherical knot $\bar{J}$ (resp. $\bar{K}$) is equivalent to J' (resp. K').

This completes the proof. □

Proof of Theorem 9.13. [8, 9] proved the following. Let V be a Seifert hypersurface for an n-dimensional closed oriented submanifold $K \subset S^{n+2}$. Then there is a Seifert hypersurface W for the $(n+2)$-submanifold $K \otimes [2] \subset S^{n+4}$ such that W is diffeomorphic to $B^{n+3} \cup B^{n+3}$ and that $B^{n+3} \cap B^{n+3}$ is diffeomorphic to $V \times [-1, 1]$.

Note that P has a Seifert hypersurface which is diffeomorphic to the punctured Poincaré sphere.

Therefore, by the above theorem in [8, 9], the $(2\nu+2)$-submanifold $P \otimes^{\nu} [2]$ has a Seifert hypersurface Q which consists of a $(2\nu+3)$-dimensional 0-handle, $(2\nu+3)$-dimensional $(\nu+1)$-handles, and $(2\nu+3)$-dimensional $(\nu+2)$-handles. Note $\nu \geqq 2$. By the van Kampen's theorem, $\pi_1(\partial Q) = 1$. By using the Meyer-Vietoris exact sequence, Q is a homology $(2\nu+3)$-ball. By the van Kampen's theorem, Q is simply-connected.

By [28], Q is diffeomorphic to the $(2\nu+3)$-ball.

Hence $P \otimes^{\nu} [2]$ has a Seifert hypersurface which is diffeomorphic to the $(2\nu+3)$-ball.

Hence $P \otimes^{\nu} [2]$ is equivalent to the trivial knot. □

14. Proof of Theorems in §10

Proof of Theorem 10.1. By Theorem 5.3, Theorem 10.1 follows from Theorem 10.2. □

Proof of Theorem 10.2. By [9], $K_* \otimes^{\nu} [2]$ has a Seifert hypersurface V_* with a handle decomposition of one $(2\nu+2)$-dimensional 0-handle and $(2\nu+2)$-dimensional $(\nu+1)$-handles, where there may not be $(2\nu+2)$-dimensional $(\nu+1)$-handles $(* = +, -, 0)$. Therefore the ν-Alexander matrix $t \cdot S_{\nu}(K_+ \otimes^{\nu} [2]) - N_{\nu}(K_+ \otimes^{\nu} [2])$ associated with V_* is 'empty'. By Proposition 12.1, $S_{\nu+1}(K_* \otimes^{\nu} [2])$ and $N_{\nu+1}(K_* \otimes^{\nu} [2])$ are square matrices. By Proposition 6.1, the $(\nu+1)$-$\mathbb{Q}[t, t^{-1}]$-Alexander polynomial of $K_* \otimes^{\nu} [2]$ is the $\mathbb{Q}[t, t^{-1}]$-balanced class of the determinant of $t \cdot S_{\nu+1}(K_* \otimes^{\nu} [2]) - N_{\nu+1}(K_* \otimes^{\nu} [2])$ associated with V_*.

By §6, $S_{\nu+1}(K_* \otimes^{\nu} [2]) = (-1)^{\nu}\, {}^tN_{\nu+1}(K_* \otimes^{\nu} [2])$.

Hence the $(\nu+1)$-Alexander matrix $P_{\nu+1}(K_* \otimes^{\nu} [2]) = (p^*_{ij})$ is equal to
$t \cdot S_{\nu+1}(K_* \otimes^{\nu} [2]) - N_{\nu+1}(K_* \otimes^{\nu} [2]) = t \cdot S_{\nu+1}(K_* \otimes^{\nu} [2]) + (-1)^{\nu+1} \cdot {}^tS_{\nu+1}(K_* \otimes^{\nu} [2])$.

By Proposition 12.1.(7), $S_{\nu+1}(K_* \otimes^{\nu} [2]) = (-1)^{\frac{\nu(\nu-1)}{2}} \quad S_1(K_*)$.

Since (K_+, K_-, K_0) is a crossing-change-triple, $S_1(K_*) = (-1)^{\frac{\nu(\nu-1)}{2}} \quad S_{\nu+1}(K_* \otimes^{\nu} [2])(* = +, -, 0)$ has the following property.

(1) $(-1)^{\frac{\nu(\nu-1)}{2}} \quad S_{\nu+1}(K_+ \otimes^{\nu} [2]) = (s^+_{ij})$ and $(-1)^{\frac{\nu(\nu-1)}{2}} \quad S_{\nu+1}(K_- \otimes^{\nu} [2]) = (s^-_{ij})$ are $\rho \times \rho$ matrices.
$(-1)^{\frac{\nu(\nu-1)}{2}} \quad S_{\nu+1}(K_0 \otimes^{\nu} [2]) = (s^0_{ij})$ is a $(\rho-1) \times (\rho-1)$ matrix $(\rho \geqq 2, \quad \rho \in \mathbb{N})$.
(2) $s^+_{\rho,\rho} - s^-_{\rho,\rho} = 1$.
(3) $s^+_{ij} = s^-_{ij} = s^0_{ij}$ $(1 \leqq i \leqq \rho-1,\ 1 \leqq j \leqq \rho-1)$. $s^+_{ij} = s^-_{ij}((i,j) \neq (\rho,\rho))$.

Note we can suppose that $\rho - 1 \geqq 1$ by using surgeries of Seifert hypersurfaces, if necessary.

$(-1)^{\frac{\nu(\nu-1)}{2}} \quad P_{\nu+1}(K_* \otimes^{\nu} [2])$ has the following properties $(* = +, -, 0)$.
(1) $(-1)^{\frac{\nu(\nu-1)}{2}} \quad P_{\nu+1}(K_+ \otimes^{\nu} [2]) = (p^+_{ij})$ and $(-1)^{\frac{\nu(\nu-1)}{2}} \quad P_{\nu+1}(K_- \otimes^{\nu} [2]) = (p^-_{ij})$ are $\rho \times \rho$ matrices.
$(-1)^{\frac{\nu(\nu-1)}{2}} \quad P_{\nu+1}(K_0 \otimes^{\nu} [2]) = (p^0_{ij})$ is a $(\rho-1) \times (\rho-1)$ matrix. $(\rho \geqq 2, \quad \rho \in \mathbb{N})$.
(2) $p^+_{\rho,\rho} - p^-_{\rho,\rho} = t + (-1)^{\nu+1}$.
(3) $p^+_{ij} = p^-_{ij} = p^0_{ij}$ $(1 \leqq i \leqq \rho-1,\ 1 \leqq j \leqq \rho-1)$. $p^+_{ij} = p^-_{ij}((i,j) \neq (\rho,\rho))$.

By calculus of determinants,

$$\mathrm{det}P_{\nu+1}(K_+ \otimes^{\nu} [2]) - \mathrm{det}P_{\nu+1}(K_- \otimes^{\nu} [2]) = c(t + (-1)^{\nu+1}) \cdot \mathrm{det}P_{\nu+1}(K_0 \otimes^{\nu} [2])$$

where $c = \pm 1$.

Hence Theorem 10.2 holds. Note that the $\mathbb{Q}[t, t^{-1}]$-Alexander polynomial is a $\mathbb{Q}[t, t^{-1}]$-balanced class. $\square$

Proof of Theorem 10.3. The 'l =even' case is proved in [23].

We prove the 'l =odd' case.

There is a Seifert hypersurface V_* for K_* ($* = +, -, 0$) such that (V_+, V_-, V_0) is related by a twist-move in B^{2l+3}.

Take Seifert matrices for K_* associated with V_* ($* = +, -, 0$). Since (V_+, V_-, V_0) is a twist-move-triple, we can suppose that $S_l(K_+) = S_l(K_-) = S_l(K_0)$ and that
$N_l(K_+) = N_l(K_-) = N_l(K_0)$.

Since K_+ is a spherical knot, the linear map which is defined by $S_l(K_+) - N_l(K_+)$ is injective. Hence the linear map which is defined by $S_l(K_*) - N_l(K_*)$ is injective ($* = +, -, 0$).

Hence the l-Alexander matrix $t \cdot S_l(K_*) - N_l(K_*)$ associated with V_* is injective. By Proposition 6.1, $A_{l+1}(K_*)$ is the determinant of the $(l + 1)$-Alexander matrix
$P_{l+1}(K_*) = t \cdot S_{l+1}(K_*) - N_{l+1}(K_*)$.

Since (K_+, K_-, K_0) is a twist-move-triple, $S_{l+1}(K_*)(* = +, -, 0)$ has the following property.
(1) $S_{l+1}(K_+) = (s^+_{ij})$ and $S_{l+1}(K_-) = (s^-_{ij})$ are $\rho \times \rho$ matrices.
$S_{l+1}(K_0) = (s^0_{ij})$ is a $(\rho - 1) \times (\rho - 1)$ matrix. ($\rho \geqq 2, \quad \rho \in \mathbb{N}$).
(2) $s^+_{\rho,\rho} - s^-_{\rho,\rho} = 1$.
(3) $s^+_{ij} = s^-_{ij} = s^0_{ij}$ ($1 \leqq i \leqq \rho - 1$, $1 \leqq j \leqq \rho - 1$). $s^+_{ij} = s^-_{ij}((i, j) \neq (\rho, \rho))$.
Note we can suppose that $\rho-1 \geqq 1$ by using surgeries of Seifert hypersurfaces, if necessary.

We have $P_{l+1}(K_*) = t \cdot S_{l+1}(K_*) - N_{l+1}(K_*) = t \cdot S_{l+1}(K_*) - (-1)^l \cdot {}^tS_{l+1}(K_*)$ Since l is even, $P_{l+1}(K_*) = t \cdot S_{l+1}(K_*) + {}^tS_{l+1}(K_*)$. $P_{l+1}(K_*)$ satisfies the following ($* = +, -, 0$).
(1) $P_{l+1}(K_+) = (p^+_{ij})$ and $P_{l+1}(K_-) = (p^-_{ij})$ are $\rho \times \rho$ matrices.
$P_{l+1}(K_0) = (p^0_{ij})$ is a $(\rho - 1) \times (\rho - 1)$ matrix ($\rho \geqq 2, \quad \rho \in \mathbb{N}$).
(2) $p^+_{\rho,\rho} - p^-_{\rho,\rho} = t + 1$.
(3) $p^+_{ij} = p^-_{ij} = p^0_{ij}$ ($1 \leqq i \leqq \rho - 1$, $1 \leqq j \leqq \rho - 1$). $p^+_{ij} = p^-_{ij}((i, j) \neq (\rho, \rho))$.

By calculus of determinants,

$$\det P_{l+1}(K_+) - \det P_{l+1}(K_-) = (t + 1) \cdot \det P_{l+1}(K_0).$$

Hence Theorem 2.11 holds. $\square$

Proof of Theorem 10.4. Since (K_+, K_-, K_0) is a twist-move-triple, there is a Seifert hypersurface V_+ (resp. V_-, V_0) for K_+(resp. K_-, K_0) such that
$S_l(K_+) = S_l(K_-) = S_l(K_0)$ and $N_l(K_+) = N_l(K_-) = N_l(K_0)$ associated with V_+, V_-, V_0.

Since K_+ is a spherical knot, we have that $S_l(K_+)$ and $N_l(K_+)$ are square matrix and furthermore we can suppose that the determinant of $S_l(K_+) - N_l(K_+)$ is ± 1. Hence we can suppose that the determinant of $S_l(K_*) - N_l(K_*)$ is ± 1 ($* = +, -, 0$).

By [9], an $(l+\nu)$-Alexander matrix $P_{l+\nu}(K_*)$ is $\pm(t\cdot S_{l+\nu}(K_*)\pm{}^tN_{l+\nu}(K_*))$. If we let $t=1$ or $t=-1$, then the determinant of $P_{l+\nu}(K_*)$ is not zero. Note that $P_{l+\nu}(K_*)$ is a square matrix because K_+ is a spherical knot. Hence the linear map which is defined by $P_{l+\nu}(K_*)$ is injective.

Hence the $(l+1+\nu)$-$\mathbb{Q}[t,t^{-1}]$-Alexander polynomial is the $\mathbb{Q}[t,t^{-1}]$-balanced class of the determinant of $P_{l+1+\nu}(K_*)$.

We have the following.

(1) $P_{l+1+\nu}(K_+\otimes^\nu[2])=(p^+_{ij})$ and $P_{l+1+\nu}(K_-\otimes^\nu[2])=(p^-_{ij})$ are $\rho\times\rho$ matrices.

$P_{l+1+\nu}(K_0\otimes^\nu[2])=(p^0_{ij})$ is a $(\rho-1)\times(\rho-1)$ matrix. $(\rho\geqq 2,\quad \rho\in\mathbb{N})$.

(2) $p^+_{\rho,\rho}-p^-_{\rho,\rho}=c(t+(-1)^{l+1+\nu})$, where $c=\pm1$.

(3) $p^+_{ij}=p^-_{ij}=p^0_{ij}$ $(1\leqq i\leqq\rho-1,\ 1\leqq j\leqq\rho-1)$. $p^+_{ij}=p^-_{ij}((i,j)\neq(\rho,\rho))$.

By calculus of determinants,

$$\det P_{l+1+\nu}(K_+\otimes^\nu[2])-\det P_{l+1+\nu}(K_-\otimes^\nu[2])=c(t+(-1)^{l+1+\nu})\cdot\det P_{l+1+\nu}(K_0\otimes^\nu[2]),$$

where $c=\pm1$.

Hence Theorem 10.4 holds. □

Proof of Theorem 10.5. By [9], $K_*\otimes^\mu$(the Hopf link) has a Seifert hypersurface V_* with a handle decomposition of one $(4\mu+2)$-dimensional 0-handle and $(4\mu+2)$-dimensional $(2\mu+1)$-handles, where there may not be $(4\mu+2)$-dimensional $(2\mu+1)$-handles $(*=+,-,0)$.

By [9], there is a Seifert matrix $S_{2\mu+1}(\,K_*\otimes^\mu$ (the Hopf link)$\,)$ associated with V_* which is equal to a Seifert matrix $(-1)^\mu S_1(K_*)$ for the 1-knot K_*. (Recall $S_*(\quad)$, $N_*(\quad)$ in §6.)

Since $K_*\otimes^\mu$ (the Hopf link) is a $(4\mu+1)$-submanifold, $N_{2\mu+1}(\,K_*\otimes^\mu$ (the Hopf link)$\,)$ $={}^tS_{2\mu+1}(\,K_*\otimes^\mu$ (the Hopf link)$\,)$. Hence we can suppose that the absolute value of the determinant of a matrix which represents the intersection product $H_{2\mu+1}(V_+;\mathbb{Z})\times H_{2\mu+1}(V_+;\mathbb{Z})\to\mathbb{Z}$ (resp. $H_{2\mu+1}(V_-;\mathbb{Z})\times H_{2\mu+1}(V_-;\mathbb{Z})\to\mathbb{Z}$) is one. Hence $K_+\otimes^\mu$ (the Hopf link) (resp. $K_-\otimes^\mu$ (the Hopf link)) is a homology sphere.

By the above handle decomposition of V_* and [28], $K_+\otimes^\mu$ (the Hopf link) (resp. $K_-\otimes^\mu$ (the Hopf link)) is PL homeomorphic to the standard sphere.

By Proposition 12.2, $(K_+\otimes^\mu$ (the Hopf link), $K_-\otimes^\mu$ (the Hopf link), $K_0\otimes^\mu$ (the Hopf link) $)$ is a twist-move-triple.

By Theorem 7.3, the proof is completed. □

Proof of Theorem 10.6. Proposition 12.1 implies the following.

$S_{p+\nu}(\,K_*\otimes^\nu[2]\,)=(-1)^cS_p(K_*)$.

$N_{p+\nu}(\,K_*\otimes^\nu[2]\,)=(-1)^{c+\nu}N_p(K_*)$. $(*=+,-,0.\quad c$ is a constant integer.)

Since K_+, K_- are spherical knots and (K_+,K_-,K_0) is a (p,q)-pass-move-triple, we have the following.

There is a $(p+\nu)$-Alexander matrix

$P_{p+\nu}(K_*\otimes^\nu[2]) = (-1)^c\{t\cdot S_p(K_*\otimes^\nu[2]) - (-1)^\nu\cdot N_p(K_*\otimes^\nu[2])\} = (p^*_{ij})$ with the following property.
(1) $P_{p+\nu}(K_+\otimes^\nu[2])$ and $P_{p+\nu}(K_-\otimes^\nu[2])$ are $\rho\times\rho$ matrices.
$P_{p+\nu}(K_0\otimes^\nu[2])$ is a $(\rho-1)\times(\rho-1)$ matrix. $(\rho \geqq 2, \quad \rho\in\mathbb{N})$
(2) $p^+_{\rho,\rho} - p^-_{\rho,\rho} = (-1)^c\{t+(-1)^{1+\nu}\}$.
(3) $p^+_{ij} = p^-_{ij} = p^0_{ij}$ $(1\leqq i\leqq \rho-1,\ 1\leqq j\leqq \rho-1)$. $p^+_{ij} = p^-_{ij}((i,j)\neq(\rho,\rho))$.

Note we can suppose that $\rho-1\geqq 1$ by using a surgery of Seifert hypersurface, if necessary.

By calculus of determinants,
$\det P_{p+\nu}(K_+\otimes^\nu[2]) - \det P_{p+\nu}(K_-\otimes^\nu[2]) = (-1)^\zeta\cdot(t+(-1)^{\nu+1})\cdot\det P_{p+\nu}(K_0\otimes^\nu[2])\cdots\cdots(!)$, where ζ is a constant integer.

Take $\det P_{p+\nu}(K_*\otimes^\nu[2])$ for each $*$ $(* = +,-,0)$. Here, there are the following three cases (i)(ii)(iii).
(i) Suppose $\det P_{p+\nu}(K_*\otimes^\nu[2])\neq 0$. Let $p-1\neq n+1-p$. Then K_+, K_- and K_0 has a same $(p-1)$-Alexander matrix $t\cdot S_{p-1}-N_{p-1}$. The $(p-1)$-Alexander matrix has the following properties. Note that $t\cdot S_{p-1}-N_{p-1}$ is a square matrix. $t\cdot S_{p-1}-N_{p-1}$ is a nonsingular square matrix. Because: K_+ and K_- are spherical knots. Hence If $t=1$, $t\cdot S_{p-1}-N_{p-1} = S_{p-1}-N_{p-1}$ is nonsingular.
By [9], a $(p+\nu-1)$-Alexander matrix $P_{p+\nu-1}$ for $K_*\otimes^\nu[2]$ is one of $\pm\{t\cdot S_{p-1}\pm N_{p-1}\}$.
If $t=1$ or $t=-1$, $P_{p+\nu-1}$ is nonsingular. Hence $P_{p+\nu-1}$ is nonsingular.
Hence the $(p+\mu)$-$\mathbb{Q}[t,t^{-1}]$-Alexander polynomial for $K_*\otimes^\nu[2]$ is $\det P_{p+\mu}(K_*\otimes^\nu[2])$.
(ii) Suppose $\det P_{p+\nu}(K_*\otimes^\nu[2])\neq 0$. Let $p-1 = n+1-p$. A $(p+\nu-1)$-Alexander matrix for $K_*\otimes^\nu[2]$ defines an injective map because a $(p-1)$-Alexander matrix for K_* does. See the identity right above Proposition 6.2.
Hence the $(p+\nu)$-$\mathbb{Q}[t,t^{-1}]$-Alexander polynomial for K_* is $\det P_{p+\nu}(K_*\otimes^\nu[2])$.
(iii) Suppose $\det P_{p+\nu}(K_*\otimes^\nu[2]) = 0$. Then the $(p+\nu)$-$\mathbb{Q}[t,t^{-1}]$-Alexander polynomial for $K_*\otimes^\nu[2]$ is 0$=\det P_{p+\nu}(K_*\otimes^\nu[2])$. Here, note that we do not need to consider whether the linear map defined by a $(p+\nu-1)$-Alexander matrix is injective or not.
By the above (i), (ii), (iii) and the identity (!) several lines above here, the proof is completed. Note that the $(p+\nu)$-$\mathbb{Q}[t,t^{-1}]$-Alexander polynomial is a $\mathbb{Q}[t,t^{-1}]$-balanced class. □

15. Discussion and open problems

We state a problem.

Problem 15.1. Suppose that n-dimensional closed oriented submanifolds K and K' differ by one twist-move (resp. pass-move). Suppose that m-submanifolds-dimensional closed oriented J and J' differ by one twist-move (resp. pass-move). Then how do we characterize a relation between $K\otimes J$ and $K'\otimes J'$?

(Recall the following. Let $n = 1$. K and K' differ by one twist-move if and only if K and K' differ by one crossing-change.)

Many variations of our results can be made.

REFERENCES

[1] J. W. Alexander: Topological invariants of knots and links, *Trans A.M.S.* 30 (1928) 275-306.

[2] E. H. Brown and B. Steer: A note on Stiefel manifolds, *Amer.J.Math.* 87 (1965) 215-217.

[3] J. Conway: An enumeration of knots and links and some of their related properties, *In:Computational problems in Abstract Algebra (Oxford 1967) Welsch Pergamon Press* (1970) 329-358.

[4] A. Haefliger: Differentiable imbeddings *Bull. Amer. Math. Soc.* 67 (1961) 109-112.

[5] A. Haefliger: Knotted (4k - 1)-Spheres in 6k-Space *Annals of Mathematics, Second Series* 75 (1962) 452-466.

[6] T. Homma: ON DEHN'S LEMMA FOR S^3 *The journal of the Yokohama Municipal University* Series D 5 (1957) 223-244.

[7] V. F. R. Jones: Hecke Algebra representations of braid groups and link *Ann. of Math.* 126 (1987) 335-388.

[8] L. H. Kauffman: Products of knots, *Bulletin of the American mathematical society* 80 (1974)1104-1107.

[9] L. H. Kauffman and W. D. Neumann: Products of knots, branched fibrations and sums of singularities, *Topology* 16 No 4 (1977) 369-393.

[10] L. H. Kauffman: On knots, *Ann of math studies* 115 (1987).

[11] K. Kawakubo: On the inertia groups of homology tori, *J.Math.Soc.Japan* 21 (1969) 37-47.

[12] M. Kervaire and J. Milnor: Groups of homotopy spheres I *Ann. of Math.* 77 (1963) 504-537.

[13] R. Kirby: The topology of 4-manifolds *Lecture Notes in Math (Springer Verlag)* vol. 1374 (1989).

[14] J. Levine: Polynomial invariant of knots of codimension two, *Ann. Math.* 84, (1966) 537-554.

[15] J. Levine: Knot cobordism in codimension two, *Comment. Math. Helv.* 44 (1969) 229-244.

[16] J. Levine: An algebraic classification of some knots of codimension two. *Comment. Math. Helv.* 45 (1970) 185–198.

[17] J.W.Milnor: Singular points of complex hyper surfaces, *Ann. of Math. Studies,no. 61, Princeton Univ. Press, Princeton, N.J.; Univ. of Tokyo Press, Tokyo,* (1968).

[18] J. W. Morgan and H. Bass eds: The Smith conjecture, *Pure and Applied Mathematics, 112* (1984).

[19] E. Ogasa: Intersectional pairs of n-knots, local moves of n-knots and invariants of n-knots, *Mathematical Research Letters* 5 (1998) 577-582, Univ. of Tokyo preprint UTMS 95-50.

[20] E. Ogasa: The intersection of spheres in a sphere and a new geometric meaning of the Arf invariants, *Journal of knot theory and its ramifications* 11 (2002) 1211-1231, Univ. of Tokyo preprint series UTMS 95-7, math.GT/0003089 in http://xxx.lanl.gov.

[21] E. Ogasa: Ribbon-moves of 2-links preserve the μ-invariant of 2-links, *Journal of knot theory and its ramifications*13 (2004) 669–687, UTMS 97-35, math.GT/0004008.

[22] E. Ogasa: Ribbon-moves of 2-knots: The Farber-Levine pairing and the Atiyah-Pathodi-Singer-Casson-Gordon-Ruberman $\tilde{\eta}$ invariant of 2-knots, *Journal of Knot Theory and Its Ramifications* 16 (2007) 523-543, math.GT/0004007, UTMS 00-22, math.GT/0407164.

[23] E. Ogasa: Local move identities for the Alexander polynomials of high-dimensional knots and inertia groups, *Journal of Knot Theory and Its Ramifications* 18 (2009) 531-545, UTMS 97-63. math.GT/0512168.

[24] E. Ogasa: A new obstruction for ribbon-moves of 2-knots: 2-knots fibred by the punctured 3-torus and 2-knots bounded by the Poicaré sphere, arXiv:1003.2473math.GT

[25] C. D. Papakyriakopoulos: On Dehn's lemma and the asphericity of knots *Proc Natl Acad Sci U S A.* 43 (1957) 169172.

[26] N. Sato: Cobordisms of semi-boundary links *Topology Appl.* 18 (1984) 225-234.

[27] M. Scharlemann: Unknotting Number One Knots Are Prime *Invent. Math.* 82 (1985) 37-55.

[28] S. Smale: Generalized Poincaré Conjecture in Dimensions Greater Than Four *Annals of Math.* 74(2) (1961) 391–406.

[29] P. A. Smith: Transformations of finite period. II, *Annals of Mathematics. Second Series* 40 (1939) 690711.

[30] H. Whitney: Differentiable Manifolds, it Annals of Mathematics Second Series 37 (1936) 645-680.

[31] H. Whitney: The Self-Intersections of a Smooth n-Manifold in 2n-Space *Annals of Mathematics Second Series* 45 (1944) 220-246.

[32] W. Whitten: Characterizations of knots and links, *Bull. of Amer. Math. Soc.* 80 (1974) 1265-1270.

[33] E, Zeeman: Twisting spun knots *Trans. Amer. Math. Soc.* 115 (1965) 471-495.

Louis H. Kauffman: Department of Mathematics, Statistics, and Computer Science, University of Illinois at Chicago, 851 South Morgan Street, Chicago, Illinois 60607-7045, USA kauffman@uic.edu

Eiji Ogasa: Computer Science, Meijigakuin University, Yokohama, Kanagawa, 244-8539, Japan pqr100pqr100@yahoo.co.jp ogasa@mail1.meijigkakuin.ac.jp